

LPTENS-93/02,
 hep-th/9302035,
 February 1993

THE QUANTUM GROUP STRUCTURE OF 2D GRAVITY AND MINIMAL MODELS II: THE GENUS-ZERO CHIRAL BOOTSTRAP

Eugène CREMMER

Jean-Loup GERVAIS

Jean-François ROUSSEL

Laboratoire de Physique Théorique de l'École Normale Supérieure¹,
 24 rue Lhomond, 75231 Paris CEDEX 05, France.

Abstract

The F and B matrices associated with Virasoro null vectors are derived in closed form by making use of the operator-approach suggested by the Liouville theory, where the quantum-group symmetry is explicit. It is found that the entries of the fusing and braiding matrices are not simply equal to quantum-group symbols, but involve additional coupling constants whose derivation is one aim of the present work. Our explicit formulae are new, to our knowledge, in spite of the numerous studies of this problem. The relationship between the quantum-group-invariant (of IRF type) and quantum-group-covariant (of vertex type) chiral operator-algebras is fully clarified, and connected with the transition to the shadow world for quantum-group symbols. The corresponding 3-j-symbol dressing is shown to reduce to the simpler transformation of Babelon and one of the author (J.-L. G.) in a suitable infinite limit defined by analytic continuation. The above two types of operators are found to coincide when applied to states with Liouville momenta going to ∞ in a suitable way. The introduction of quantum-group-covariant operators in the three dimensional picture gives a generalisation of the quantum-group version of discrete three-dimensional gravity that includes tetrahedra associated with 3-j symbols and universal R-matrix elements. Altogether the present work gives the concrete realization of Moore and Seiberg's scheme that describes the chiral operator-algebra of two-dimensional gravity and minimal models.

¹Unité Propre du Centre National de la Recherche Scientifique, associée à l'École Normale Supérieure et à l'Université de Paris-Sud.

1 INTRODUCTION

Clearly, the chiral operator-algebras of two dimensional (2D) gravity, and minimal models have already received much attention. It may thus be surprising that the corresponding chiral operator algebra is not yet fully explored. There are several reasons for this situation.

In recent years the emphasis has been put on what is considered as particularly simple and elegant in conformal theories. On the one hand, Verlinde's beautiful ideas[1] led to extensive studies of the fusion rules where one only considers the number of independent couplings –multiplicities– N_{ij}^k which are non-negative integers. One aim here is, on the contrary, to study the fusion algebra, that is to determine the coefficients of Wilson's operator-product-expansion OPE completely (since we deal with irrational theories, the N_{ij}^k are trivially known)². On the other hand, after the topological nature of the gravity-dressed minimal models was recognized, and since the recent developments are connected with matrix models, most recent articles deal with topological conformal theories, where the chiral OPE we want to study becomes somewhat hidden.

Most previous studies of the operator-product algebra (OPA) focused on the braiding matrices[2, 3, 4, 5, 6, 7, 8], a few years after ref.[9] first determined the simplest ones. Fusion was only used to leading order as a tool to deduce the higher braiding matrices from the simplest ones, recursively.

One main result of our study is the explicit correspondence between q-6j symbols and the fusing (F) and braiding (B) matrices of the basic set of chiral vertex operators $V_m^{(J)}$ of the type $(1, 2J + 1)$ in the BPZ classification, with $2J$ a positive integer. In the notation of Moore and Seiberg (MS)[10], we will show that

$$F_{J_{23}, J_{12}} \begin{bmatrix} J_1 & J_2 \\ J & J_3 \end{bmatrix} = \frac{g_{J_1 J_2}^{J_{12}} g_{J_{12} J_3}^J}{g_{J_2 J_3}^{J_{23}} g_{J_1 J_{23}}^J} \left\{ \begin{matrix} J_1 & J_2 \\ J_3 & J \end{matrix} \middle| \begin{matrix} J_{12} \\ J_{23} \end{matrix} \right\}. \quad (1.1)$$

The F and B matrices are found to be connected by the MS relation[10] (the Δ 's are conformal weights)

$$B_{J_{23}, J_{12}}^\pm \begin{bmatrix} J_1 & J_3 \\ J & J_2 \end{bmatrix} = e^{\pm i\pi(\Delta_J + \Delta_{J_2} - \Delta_{J_{23}} - \Delta_{J_{12}})} F_{J_{23}, J_{12}} \begin{bmatrix} J_1 & J_2 \\ J & J_3 \end{bmatrix}. \quad (1.2)$$

In the first equation, $\left\{ \begin{matrix} J_1 & J_2 \\ J_3 & J \end{matrix} \middle| \begin{matrix} J_{12} \\ J_{23} \end{matrix} \right\}$ is the q-6j symbol associated with $U_q(sl(2))$, and we have introduced quantities noted g_{JK}^L , whose explicit expression will be given in section 2. The V operators are closely related with operators called IRF-chiral vertex operator in ref.[11], which are associated with integrable models with solid-on-solid interactions around-the-face[12]. In the context of Liouville theory, they were called[13] Bloch-wave operators since they diagonalise the monodromy matrix, and we shall use this name. The above formula is new to our knowledge, although many other studies[3, 5, 6, 11, 14, 15, 16, 17, 18] gave strong hints that indeed the fusion and braiding matrices of the chiral vertex operators should be “essentially given” by q-6j symbols. The basic difficulty in trying to relate OPA coefficients with quantum-group symbols is that the latter are trigonometric functions of the quantum-group parameters, while the correlators of Bloch-wave operators involve ordinary Γ functions. Thus equality is impossible, although, both satisfy the same MS polynomial

²Warning! throughout the article the word fusion is related with full-fledged OPE, not with Verlinde algebras

equations. The form of the above formulae is precisely such that this crucial property holds. An important remark must be made at this point. We have derived Eq.1.1 with a specific normalization of the Bloch-wave operators, that is, that their matrix elements between highest-weight states are equal to one. Since it leaves invariant the polynomial equations, the multiplicative factor $g_{J_1 J_2}^{J_{12}} g_{J_{12} J_3}^J / g_{J_2 J_3}^{J_{23}} g_{J_1 J_{23}}^J$ is a gauge transformation, from the three dimensional viewpoint, which may be re-absorbed by a change of this normalization. However this does not eliminate the g 's from the OPE, as one may easily see, for instance at the level of primaries. Indeed, the fusing-matrix entries are equal to the coefficients of the OPA for primaries only if the vertex operators are normalized to one (this will be spelled out later on). Accordingly, the difference between our result and 6-j symbols is genuine. Returning to our main line, let us first sketch how the operator algebra of the V fields is determined in section 2. At first $F_{J_{23}, J_{12}} \begin{bmatrix} 1/2 & J_2 \\ J & J_3 \end{bmatrix}$ is re-derived from the hypergeometric differential equation due to the decoupling of the Virasoro null vector at the second level. Next, the result is put under the form Eq.1.1. Remarkably, this determines the coupling constants g_{JK}^L uniquely, assuming, as is always possible that $g_{JK}^{J+K} = 1$. Then the complete operator-algebra is established applying Moore and Seiberg's general scheme.

Another difficulty of the quantum-group picture is that formulae like Eqs.1.1, 1.2 only involve quantum-group invariants, so that one does not see how the quantum group acts. Previously, this problem has been solved in two seemingly different ways. First the invariant operators were “dressed with 3-j symbols” [12, 11]. In this approach, the quantum group covariance seems somewhat artificial and redundant, even though this construction allows[12] to relate integrable models. of the IRF (interaction around-the-face) and vertex types. Another method[19, 13, 30, 20] was directly inspired by the operator approach[21, 9, 22, 23] to Liouville theory which is explicitly quantum-group symmetric. A set of chiral primary fields noted $\xi_M^{(J)}$ was constructed³. They are deduced from the Bloch-wave operators by equations of the form[13, 19]

$$\xi_M^{(J)} := \sum_{-J \leq m \leq J} |J, \varpi\rangle_M^m E_m^{(J)}(\varpi) V_m^{(J)}, \quad -J \leq M \leq J, \quad (1.3)$$

where $|J, \varpi\rangle_M^m$ are q-hypergeometrix functions of $e^{ih(\varpi+m)}$, $E_m^{(J)}(\varpi)$ are normalization factors, and ϖ is the zero-mode. The braiding matrices of the ξ 's coincide[13, 19] with the universal R-matrix of $U_q(sl(2))$. This construction is more economical than the dressing by 3-j symbols, since it does not involve any redundant quantum number. Thus we call it the *intrinsic transformation*. The leading-order fusion coefficients of the ξ fields were shown[19] to coincide with the quantum 3-j symbols, and it was stated[20] without proof that this is also true, up to a coupling constant, for every order. It is another purpose of the present work to complete that picture. At first, using the above relationship between ξ fields and V fields, we deduce, in section 3, that the fusion of the former are given by

$$\xi_{M_1}^{(J_1)} \xi_{M_2}^{(J_2)} = \sum_{J_{12}=|J_1-J_2|}^{J_1+J_2} g_{J_1 J_2}^{J_{12}}(J_1, M_1; J_2, M_2 | J_{12}) \times$$

³There exists a related derivation of the universal R matrix in the Coulomb gas picture[24]. However it seems not to be so well suited for obtaining the complete fusion algebra.

$$\sum_{\{\nu\}} \xi_{M_1+M_2}^{(J_{12}, \{\nu\})} < \varpi_{J_{12}}, \{\nu\} | V_{J_2-J_{12}}^{(J_1)} | \varpi_{J_2} >, \quad (1.4)$$

where $(J_1, M_1; J_2, M_2 | J_{12})$ are the q-Clebsch-Gordan coefficients. Apart from the fact that the right-hand side involves one ξ field and one V field, this has the standard MS form⁴ (worldsheet variables are suppressed, the states $|\varpi_J, \{\nu\}\rangle$ span the Verma module with highest weight state $|\varpi_J\rangle$). It realizes our expectation that there must be fields such that the fusion corresponds to making q-tensor-products of representations. However there is the additional factor $g_{J_1 J_2}^{J_{12}}$. Since this form may be considered as governed by the q-deformed Wigner Eckart theorem, the g 's will be called coupling constants. It is thus clear that the ξ fields are quantum-group covariant. Next we use the fact that the right-hand side involves one ξ field and one V field to relate them operatorially. This is found equivalent to the dressing by 3-j symbols of refs.[12, 11]. Since the ξ fields were defined by the intrinsic transformation[13, 19] (see Eq.1.3), we are able to establish the relationship between the two viewpoints: the dressing by 3-j symbols has an additional magnetic quantum number, and is shown to reduce to the intrinsic transformation, when this number tends to ∞ after a suitable analytic continuation. Finally, we terminate section 3, by giving a general formula connecting the coupling constants g_{JK}^L with quantities introduced earlier[19, 20] in the quantum-group context.

In section 4, we further develop the idea of understanding the connection between V and ξ fields from infinite limits. Inspired by a recent article of Witten[15] we show that the V and ξ fields coincide in the limit where the zero-mode ϖ goes to ∞ , after suitable analytic continuation. This is explicitly proven from the intrinsic transformation summarized above (Eq.1.3). The expressions of the fusing and braiding matrices of V and ξ fields in terms of q symbols then show that these symbols should be related by the same limit, and this is explicitly verified. This sheds light on the method followed in ref[15] to construct covariant vertex operators, although the conformal theory considered is different. Another point of this section is to show that the intrinsic transformation is the conformal analogue of the transition to the shadow world described in ref.[25] for quantum group symbols, and to display its three-dimensional aspect where 3-j symbols and R-matrix are represented by tetrahedra, at the same time as 6-j symbols.

Finally, in section 5 we take account of the fact that, for irrational theories, there are actually two quantum-group parameters, one for each screening charge. The above picture should thus be extended. To avoid lengthy discussion, we take a short cut and only determine the coupling constants, for which a general formula is written down. The underlying quantum group structure was already unravelled before in refs.[13, 20, 26].

2 THE BRAIDING AND FUSING MATRICES

The present operator algebra may, in principle be considered on any given Riemann surface. In a typical situation, one may work on the cylinder $0 \leq \sigma \leq 2\pi$, $-\infty \leq \tau \leq \infty$ obtained by an appropriate mapping from one of the handles. However, the present work deals with the case of genus zero only. In this discussion we concentrate on the holomorphic fields which are functions of $x = \sigma - i\tau$ only. The notations are

⁴It already appears, without the g factor in ref.[11].

exactly the same as in refs.[19, 20, 27]. The basic concepts are recalled in appendix A for completeness. The starting point is the differential equation[21, 19], satisfied by the two spin 1/2 fields $V_{\pm 1/2}(x)$, which is recalled in appendix A (Eq.A.8). In the BPZ framework it expresses the vanishing of the Virasoro null fields at the second level for operators of the type $(1, 2)$ (Until section 5 we concentrate on operators of the $(1, 2J + 1)$ type, that is on the family with one of the two quantum group parameters). It then follows that for any holomorphic primary operator $A_{\Delta}(x)$ with conformal weight Δ , one has (see Eq.A.6 of ref[19]):

$$\begin{aligned} < \varpi_4 | V_{\pm 1/2}(x) A_{\Delta}(x') | \varpi_1 > = e^{ix'(\varpi_1^2 - \varpi_4^2)h/4\pi} e^{i(x'-x)(-1/2 \mp \varpi_4)h/2\pi} \\ & \times (1 - e^{-i(x-x')})^{\beta} F(a_{\pm}, b_{\pm}; c_{\pm}; e^{i(x'-x)}) \end{aligned} \quad (2.1)$$

$$\begin{aligned} a_{\pm} &= \beta - \frac{h}{2\pi} \mp h(\frac{\varpi_4 - \varpi_1}{2\pi}); \quad b_{\pm} = \beta - \frac{h}{2\pi} \mp h(\frac{\varpi_4 + \varpi_1}{2\pi}); \quad c_{\pm} = 1 \mp \frac{h\varpi_4}{\pi}; \\ \beta &= \frac{1}{2}(1 + h/\pi)(1 - \sqrt{1 - \frac{8h\Delta}{2\pi(1 + h/\pi)^2}}); \end{aligned} \quad (2.2)$$

where $F(a, b; c; z)$ is the standard hypergeometric function. Similarly, one verifies that

$$\begin{aligned} < \varpi_4 | A_{\Delta}(x') V_{\pm 1/2}(x) | \varpi_1 > = e^{ix'(\varpi_1^2 - \varpi_4^2)h/4\pi} e^{i(x-x')(-1/2 \pm \varpi_1)h/2\pi} \times \\ & (1 - e^{-i(x-x')})^{\beta} F(a'_{\pm}, b'_{\pm}; c'_{\pm}; e^{-i(x-x')}) \end{aligned} \quad (2.3)$$

$$a'_{\pm} = \beta - \frac{h}{2\pi} \mp h(\frac{\varpi_1 - \varpi_4}{2\pi}); \quad b'_{\pm} = \beta - \frac{h}{2\pi} \mp h(\frac{\varpi_1 + \varpi_4}{2\pi}); \quad c'_{\pm} = 1 \mp \frac{h\varpi_1}{\pi}. \quad (2.4)$$

As already mentioned in the introduction, ϖ denotes the zero mode, and $|\varpi >$ is the corresponding highest-weight state. The two fields $V_{\pm 1/2}$ have the same conformal weight, but shift ϖ in opposite directions. These are the basic equations for establishing the operator algebra. They were already used[19] to determine the operator-product algebra (OPA) to leading order in the singularity. This showed that there are operators $V_m^{(J)}$, with $2J$ a positive integer, and $-J \leq m \leq J$, (also denoted $V^{\mu, \nu}$, with $2J = \mu + \nu$, and $2m = -\mu + \nu$) and conformal weights

$$\Delta_J = -hJ(J + 1)/\pi - J. \quad (2.5)$$

The above formulae were also used in order to determine the braiding properties of the V operators. We shall only deal with the case of genus zero, so that it is appropriate, following ref.[27] to change the normalization of the operator. In this connection, let us recall that a primary field $A(z, z^*)$ of weight Δ , $\bar{\Delta}$ transforms so that $A(z, z^*)(dz)^{\Delta}(dz^*)^{\bar{\Delta}}$ is invariant[31]. Given $A(z, z^*)$, and a conformal map $z \rightarrow Z(z)$, it is thus convenient to define

$$A(Z, Z^*) = \left(\frac{dz}{dZ}\right)^{\Delta} \left(\frac{dz^*}{dZ^*}\right)^{\bar{\Delta}} U_{\{Z\}} A(z, z^*) U_{\{Z\}}^{-1}, \quad (2.6)$$

where $U_{\{Z\}}$ is the operator that realizes the transformation of the states on which A acts. It is a functional of $Z(z)$. We are using coordinates such that the Riemann sphere is described by the complex variable $z = \exp(ix)$. If we denote by $A(\sigma, \tau)$ a general primary field on the cylinder, and by $A(z, z^*)$ its transformed on the sphere, one has

$$A(z, z^*) = z^{-\Delta} z^{*- \bar{\Delta}} U_{\{z\}} A(\sigma, \tau) U_{\{z\}}^{-1}. \quad (2.7)$$

$A(\sigma, \tau)$, which was used before ref[27], is such that its Fourier expansion in σ has integer coefficients. The transformation rule just recalled gives the standard definition for the Laurent expansion of $A(z, z^*)$ as series in $z^{n-\Delta}$, and $(z^*)^{m-\Delta}$, n, m integers⁵. Using the formulae just recalled, one may see that

$$\begin{aligned} < \varpi_4 | V_{\pm 1/2}^{(1/2)}(z_3) V_{m_2}^{(J_2)}(z_2) | \varpi_1 > = z_2^{-\Delta_{J_2} - \Delta(\varpi_1)} z_3^{-\Delta_{1/2} + \Delta(\varpi_4)} \times \\ & \left(\frac{z_2}{z_3} \right)^{\Delta(\varpi_4 \pm 1)} \left(1 - \frac{z_2}{z_3} \right)^{-hJ_2/\pi} F(a_{\pm}, b_{\pm}; c_{\pm}; \frac{z_2}{z_3}); \\ a_{\pm} &= \frac{h}{2\pi} [-1 - 2J_2 \mp (\varpi_4 - \varpi_1)]; \quad b_{\pm} = \frac{h}{2\pi} [-1 - 2J_2 \mp (\varpi_4 + \varpi_1)]; \\ c_{\pm} &= 1 \mp \frac{h\varpi_4}{\pi}. \end{aligned} \quad (2.8)$$

The symbols $|\varpi >$ represent highest-weight states, with Virasoro weights $\Delta(\varpi)$ given by Eq.A.11. According to Eqs.A.12, $V_m^{(J)}$ shifts ϖ by $2m$, thus $\varpi_1 = \varpi_4 + 2m_2 \pm 1$. In order to avoid clumsy formulae, we do not explicitly write down the operators $U_{\{z\}}$, and $U_{\{z\}}^{-1}$ that appear when Eq.2.7 is used. There should be no confusion. One may consider Eq.2.8 for arbitrary ϖ_1 . However, in the framework of standard conformal theory, a basic assumption is that all highest-weight states are generated by applying a primary field to the $sl(2, C)$ invariant vacuum. As discussed, in ref.[27], a highest-weight state $|\varpi_J >$, is created from the $sl(2, C)$ -invariant state $|\varpi_0 >$ ($\varpi_0 = 1 + \pi/h$) by the limit

$$|\varpi_J > = \lim_{z \rightarrow 0} V_{-J}^{(J)}(z) |\varpi_0 >, \quad \text{with } \varpi_J = \varpi_0 + 2J. \quad (2.10)$$

We shall thus consider matrix elements between states with momenta of this type. According to Eq.A.17, the overall normalization of Eq.2.1, 2.3, and 2.8 is fixed by assuming that the operators $V_m^{(J)}$ are such that

$$< \varpi_{J_1} | V_m^{(J_2)}(1) | \varpi_{J_3} > = \delta_{m, J_3 - J_1}. \quad (2.11)$$

Similarly, one also deduces from Eq.2.3 that

$$\begin{aligned} < \varpi_4 | V_{m_2}^{(J_2)}(z_2) V_{\pm 1/2}^{(1/2)}(z_3) | \varpi_1 > = z_2^{-\Delta_{J_2} + \Delta(\varpi_4)} z_3^{-\Delta_{1/2} - \Delta(\varpi_1)} \times \\ & \left(\frac{z_2}{z_3} \right)^{\Delta(\varpi_1 \mp 1)} \left(1 - \frac{z_2}{z_3} \right)^{-hJ_2/\pi} F(a'_{\pm}, b'_{\pm}; c'_{\pm}; \frac{z_3}{z_2}) \\ a'_{\pm} &= \frac{h}{2\pi} [-1 - 2J_2 \mp (\varpi_1 - \varpi_4)]; \quad b'_{\pm} = \frac{h}{2\pi} [-1 - 2J_2 \mp (\varpi_1 + \varpi_4)]; \\ c'_{\pm} &= 1 \mp \frac{h\varpi_1}{\pi}. \end{aligned} \quad (2.12)$$

Next using the well known relation

$$F(a, b; c; x) = \frac{\Gamma(c)\Gamma(c-b-a)}{\Gamma(c-a)\Gamma(c-b)} F(a, b; a+b-c+1; 1-x) +$$

⁵The main practical advantage of using z, z^* is that $A(z, z^*)$ has a simple behaviour under Möbius transformations

$$\frac{\Gamma(c)\Gamma(a+b-c)}{\Gamma(a)\Gamma(b)} (1-x)^{c-a-b} F(c-a, c-b; c-a-b+1; 1-x), \quad (2.14)$$

one deduces that

$$\begin{aligned} & \langle \varpi_4 | V_{\pm 1/2}^{(1/2)}(z_3) V_{m_2}^{(J_2)}(z_2) | \varpi_1 \rangle = z_2^{2(\Delta_{1/2} + \Delta(\varpi_4) - \Delta(\varpi_1))} z_3^{-2\Delta_{1/2}} \times \\ & \sum_{\epsilon=\pm 1} f_{\pm 1, \epsilon}(J_2, m_2; \varpi) \langle \varpi_1 | V_{m_4}^{(J_4)}(z_2) V_{-\epsilon/2}^{(1/2)}(z_2(1 - z_2/z_3)) | \varpi_2 \rangle, \end{aligned} \quad (2.15)$$

where $m_4 = (\varpi_2 - \varpi_1 - \epsilon)/2$, and

$$\begin{aligned} f_{1,1} &= \frac{\Gamma(1 - \varpi_4 h/\pi) \Gamma(1 + (2J_2 + 1) h/\pi)}{\Gamma(1 + (-\varpi_4 + J_2 - m_2) h/\pi) \Gamma(1 + (J_2 + m_2 + 1) h/\pi)}, \\ f_{-1,-1} &= \frac{\Gamma(1 + \varpi_4 h/\pi) \Gamma(-1 - (2J_2 + 1) h/\pi)}{\Gamma(-(J_2 + m_2) h/\pi) \Gamma((\varpi_4 - J_2 + m_2 - 1) h/\pi)}, \\ f_{1,-1} &= \frac{\Gamma(1 - \varpi_4 h/\pi) \Gamma(-1 - (2J_2 + 1) h/\pi)}{\Gamma(-(J_2 - m_2) h/\pi) \Gamma(-(\varpi_4 + J_2 + m_2 + 1) h/\pi)}, \\ f_{-1,1} &= \frac{\Gamma(1 + \varpi_4 h/\pi) \Gamma(1 + (2J_2 + 1) h/\pi)}{\Gamma(1 + (\varpi_4 + J_2 + m_2) h/\pi) \Gamma(1 + (J_2 - m_2 + 1) h/\pi)}. \end{aligned} \quad (2.16)$$

Following ref.[10] we re-transform the right-hand side of Eq.2.15, in order to see the operator-product expansion more clearly. Denote by $|\varpi, \{\nu\}\rangle$ an arbitrary vector of the Verma module corresponding to the highest weight $\Delta(\varpi)$. The notation $\{\nu\}$ represents a multi-index. We shall use a basis such that the metric of inner products $\mathcal{G}_{\{\nu\}, \{\nu'\}}$ is equal to $\delta_{\{\nu\}, \{\nu'\}}$. It is convenient to write

$$\begin{aligned} & \langle \varpi_1 | V_{m_4}^{(J_4)}(z_2) V_{-\epsilon/2}^{(1/2)}(z_2(1 - z_2/z_3)) | \varpi_2 \rangle = \\ & \sum_{\{\nu\}} \langle \varpi_1 | V_{m_4}^{(J_4)}(z_2) | \varpi_\epsilon, \{\nu\} \rangle \langle \varpi_\epsilon, \{\nu'\} | V_{-\epsilon/2}^{(1/2)}\left(z_2(1 - \frac{z_2}{z_3})\right) | \varpi_2 \rangle, \end{aligned} \quad (2.17)$$

where $\varpi_\epsilon = \varpi_2 + \epsilon$. According to Eq.2.11, one has

$$\langle \varpi_1 | V_{m_4}^{(J_4)}(z_2) | \varpi_\epsilon, \{0\} \rangle \equiv \langle \varpi_1 | V_{m_4}^{(J_4)}(z_2) | \varpi_\epsilon \rangle = z_2^{\Delta_1 + \Delta_4 - \Delta_\epsilon}, \quad (2.18)$$

where

$$\Delta_1 = \Delta(\varpi_1), \quad \Delta_4 = \Delta(\varpi_4), \quad \Delta_\epsilon = \Delta(\varpi_\epsilon). \quad (2.19)$$

On the other hand,

$$\langle \varpi_4 | V_{m_2 \pm 1/2}^{(J_\epsilon)}(z_2) | \varpi_1 \rangle = z_2^{-\Delta_1 + \Delta_4 + \Delta_\epsilon}, \quad (2.20)$$

and we get

$$\langle \varpi_1 | V_{m_4}^{(J_4)}(z_2) | \varpi_\epsilon \rangle = z_2^{2(\Delta_1 - \Delta_4)} \langle \varpi_4 | V_{m_2 \pm 1/2}^{(J_\epsilon)}(z_2) | \varpi_1 \rangle.$$

It is thus consistent to let

$$\langle \varpi_1 | V_{m_4}^{(J_4)}(z_2) | \varpi_\epsilon, \{\nu\} \rangle = z_2^{2(\Delta_1 - \Delta_4)} \langle \varpi_4 | V_{m_2 \pm 1/2}^{(J_\epsilon, \{\nu\})}(z_2) | \varpi_1 \rangle, \quad (2.21)$$

where the right-hand side involves the descendant fields, noted $V_m^{(J, \{\nu\})}$ in general. This expresses the symmetry between the three legs (sphere with three punctures). The last equation allows us to re-write Eq.2.15 as

$$\begin{aligned} < \varpi_4 | V_{\pm 1/2}^{(1/2)}(z_3) V_{m_2}^{(J_2)}(z_2) | \varpi_1 > = \left(\frac{z_2}{z_3} \right)^{2\Delta_{1/2}} \sum_{\epsilon=\pm 1} f_{\pm 1, \epsilon}(J_2, m_2; \varpi_4) \times \\ \sum_{\{\nu\}} < \varpi_4 | V_{m_2 \pm 1/2}^{(J_\epsilon, \{\nu\})}(z_2) | \varpi_1 > < \varpi_\epsilon, \{\nu\} | V_{-\epsilon/2}^{(1/2)} \left(z_2 \left(1 - \frac{z_2}{z_3} \right) \right) | \varpi_2 >. \end{aligned} \quad (2.22)$$

This formula is not explicitly invariant under z translation. This apparent contradiction is resolved in appendix B, by performing a suitable Möbius transformation. At this point it is useful to establish a closer contact with the Moore Seiberg (MS) scheme[10]. There is a Verma module $\mathcal{H}_J$ for each ϖ_J (defined in Eq.2.10). The MS chiral vertex-operators connect three specified Verma modules and are thus of the form $\phi_{J_3, J_1}^{J_2}$. The Bloch-wave operators $V_m^{(J)}$, on the contrary, act in the direct sum $\mathcal{H} = \oplus_J \mathcal{H}_J$. According to Eq.2.11, the two are related by the projection operator $\mathcal{P}_J$:

$$\mathcal{P}_J \mathcal{H} = \mathcal{H}_J, \quad \mathcal{P}_{J_3} V_m^{(J_2)} \equiv \phi_{J_3, J_3+m}^{J_2}. \quad (2.23)$$

In the MS formulation, the fusing and braiding algebras of the V operators, including the explicit contribution of the descendants, read (from now on we omit the worldsheet variables):

$$\begin{aligned} \mathcal{P}_J V_{m_1}^{(J_1)} V_{m_2}^{(J_2)} &= \mathcal{P}_J \sum_{J_{12}=|J_1-J_2|}^{J_1+J_2} F_{J+m_1, J_{12}} \begin{bmatrix} J_1 & J_2 \\ J & J+m_1+m_2 \end{bmatrix} \times \\ &\sum_{\{\nu_{12}\}} V_{m_1+m_2}^{(J_{12}, \{\nu_{12}\})} < \varpi_{J_{12}}, \{\nu_{12}\} | V_{J_2-J_{12}}^{(J_1)} | \varpi_{J_2} >, \end{aligned} \quad (2.24)$$

$$\mathcal{P}_J V_{m_1}^{(J_1)} V_{m_2}^{(J_2)} = \mathcal{P}_J \sum_{m'_1, m'_2, m'_1+m'_2=m_1+m_2} B_{J+m_1, J+m'_2}^{\pm} \begin{bmatrix} J_1 & J_2 \\ J & J+m_1+m_2 \end{bmatrix} V_{m'_2}^{(J_2)} V_{m'_1}^{(J_1)}. \quad (2.25)$$

As already recalled in the introduction, the F and B matrices should be related by the equations [10]

$$B_{J_{23}, J_{12}}^{\pm} \begin{bmatrix} J_1 & J_3 \\ J & J_2 \end{bmatrix} = e^{\pm i\pi(\Delta_J + \Delta_{J_2} - \Delta_{J_{23}} - \Delta_{J_{12}})} F_{J_{23}, J_{12}} \begin{bmatrix} J_1 & J_2 \\ J & J_3 \end{bmatrix}. \quad (2.26)$$

Comparing Eq.2.24 with Eq.2.22, one concludes that⁶

$$\begin{aligned} F_{J+\epsilon_1/2, J_2+\epsilon_2/2} \begin{bmatrix} 1/2 & J_2 \\ J & J_3 \end{bmatrix} &= f_{\epsilon_1, \epsilon_2}(J_2, J_3 - J - \epsilon_1/2; \varpi_J) = \\ &\frac{\Gamma((1-\epsilon_1) - \epsilon_1(2J+1)h/\pi)}{\Gamma(1 + (\epsilon_2 - \epsilon_1)/2 + (J_3 + \epsilon_2 J_2 - \epsilon_1 J + (1 - \epsilon_1 + \epsilon_2)/2)h/\pi)} \times \\ &\frac{\Gamma(\epsilon_2 + \epsilon_2(2J_2+1)h/\pi)}{\Gamma((\epsilon_2 - \epsilon_1)/2 + (-J_3 + \epsilon_2 J_2 - \epsilon_1 J + (-1 - \epsilon_1 + \epsilon_2)/2)h/\pi)}. \end{aligned} \quad (2.27)$$

Using the explicit form of the braiding algebra of ref.[9] (or ref.[19]), one easily checks that Eq.2.26 holds for $J_1 = 1/2$.

⁶we slightly change the choice of indices for later convenience

Next we want to deduce the complete F and B matrices from the ones just written using the polynomial equations[10] that express the associativity of the OPA, and the fact that the order between fusing and braiding should be irrelevant. It is well known that the q -6j symbols with appropriate entries give a solution. However they only involve trigonometric functions of h , while the above partial solution Eq.2.27 involves Γ functions. At this point we remark that, given a solution of the polynomial equations, one may get a family of other solutions if one multiplies by factors whose products is equal to one in every equation. With this in mind, we consider the ansatz

$$F_{J_{23}, J_{12}} \begin{bmatrix} J_1 & J_2 \\ J & J_3 \end{bmatrix} = \frac{g_{J_1 J_2}^{J_{12}} g_{J_{12} J_3}^J}{g_{J_{23}}^{J_{12}} g_{J_1 J_{23}}^J} \left\{ \begin{matrix} J_1 & J_2 \\ J_3 & J \end{matrix} \middle| \begin{matrix} J_{12} \\ J_{23} \end{matrix} \right\}. \quad (2.28)$$

The symbol⁷ $\left\{ \begin{matrix} J_1 & J_2 \\ J_3 & J \end{matrix} \middle| \begin{matrix} J_{12} \\ J_{23} \end{matrix} \right\}$ represents the 6-j coefficient which is not completely tetrahedron-symmetric, as the notation indicates. This is in contrast with the Racah-Wigner 6-j symbol, noted $\left\{ \begin{matrix} J_1 & J_2 & J_{12} \\ J_3 & J & J_{23} \end{matrix} \right\}$. The form of the gg/gg -factor is precisely such that the left-hand side is also a solution of the polynomial equations, (if the B matrix is defined by Eq.2.26) for arbitrary g 's. We will give a very simple proof of this fact later on, following the argument indicated in the introduction. For the time being, let us show that the consistency of the ansatz just written, with the value of F for $J_1 = 1/2$ (Eq.2.27) fixes g_{JK}^L for arbitrary J, K, L so that F matrix is completely determined. This is of course due to the very special form of the proportionality factor which is dictated by the polynomial equations (below we shall see how the polynomial equations arise from associativity of the OPA in our case). In order to determine the g 's, we introduce

$$A \begin{bmatrix} J_1 & J_2 & J_{12} \\ J_3 & J & J_{23} \end{bmatrix} \equiv \frac{F_{J_{23}, J_{12}} \begin{bmatrix} J_1 & J_2 \\ J & J_3 \end{bmatrix}}{\left\{ \begin{matrix} J_1 & J_2 \\ J_3 & J \end{matrix} \middle| \begin{matrix} J_{12} \\ J_{23} \end{matrix} \right\}}. \quad (2.29)$$

Next we determine $A \begin{bmatrix} 1/2 & J_2 & J_{12} \\ J_3 & J & J_{23} \end{bmatrix}$ by using Eq.2.27, together with the explicit expression of the 6-j symbols (see, e.g. ref.[25] and Eq.2.55 below). After some calculations, one obtains

$$A \begin{bmatrix} 1/2 & J_2 & J_{2+\epsilon_2/2} \\ J_3 & J & J_{J+\epsilon_1/2} \end{bmatrix} = \sqrt{\frac{F(\epsilon_2 + \epsilon_2(2J_2 + 1)h/\pi)}{F((1 + (\epsilon_2 - \epsilon_1)/2) + (\epsilon_2 J_2 - \epsilon_1 J + J_3 + (1 - \epsilon_1 + \epsilon_2)/2)h/\pi)}} \\ \times \sqrt{\frac{F((1 - \epsilon_1) - \epsilon_1(2J + 1)h/\pi)}{F((\epsilon_2 - \epsilon_1)/2 + (\epsilon_2 J_2 - \epsilon_1 J - J_3 + (-1 - \epsilon_1 + \epsilon_2)/2)h/\pi)}}, \quad (2.30)$$

$$\text{with} \quad F(z) \equiv \Gamma(z)/\Gamma(1 - z). \quad (2.31)$$

The obvious property $F(z)F(1 - z) = 1$ is important for the following. At this point our ansatz gives

$$\frac{g_{J_2+\epsilon_2/2, J_3}^J}{g_{J_2 J_3}^{J+\epsilon_1/2}} = \frac{g_{1/2, J+\epsilon_1/2}^J}{g_{1/2, J_2}^{J+\epsilon_2/2}} A \begin{bmatrix} 1/2 & J_2 & J_{2+\epsilon_2/2} \\ J_3 & J & J_{J+\epsilon_1/2} \end{bmatrix}. \quad (2.32)$$

Clearly, once we know the expression of $g_{1/2 J}^{J\pm 1/2}$, this last relation will allow us to determine all of the $g_{J_1 J_2}^{J_{12}}$ by a double recursion on the indices J_1 and J_{12} (for any J_2).

⁷since all symbols are q deformed, we omit the letter q most of the time.

The next basic point is that the integrability condition for this recurrence relation, completely fixes $g_{1/2 J}^{J+1/2}$, as we now show. As a preliminary remark, note that $g_{1/2 J}^{J+1/2}$ can be taken equal to one, without loss of generality. For, if it were not, it would be possible to define

$$\tilde{g}_{J_1 J_2}^{J_{12}} \equiv \frac{\alpha_{J_1} \alpha_{J_2}}{\alpha_{J_{12}}} g_{J_1 J_2}^{J_{12}} \quad \text{such that} \quad \tilde{g}_{1/2 J}^{J+1/2} \equiv \frac{\alpha_{1/2} \alpha_J}{\alpha_{J+1/2}} g_{1/2 J}^{J+1/2} = 1,$$

and the fusion coefficients F defined by Eq.2.28 with $\tilde{g}$ are the same as the ones defined with g . Returning to our main line, it is useful to consider Eq.2.32 as defining a connection on a two dimensional lattice, for fixed J_3 , by letting

$$U_{-\epsilon_1, \epsilon_2}^{(J_3)}(J + \frac{\epsilon_1}{2}, J_2) \equiv \frac{g_{1/2, J+\epsilon_1/2}^J}{g_{1/2, J_2}^{J_2+\epsilon_2/2}} A \left[\begin{matrix} 1/2 & J_2 & J_2+\epsilon_2/2 \\ J_3 & J & J+\epsilon_1/2 \end{matrix} \right]. \quad (2.33)$$

This connection is flat, since Eq.2.32 is equivalent to

$$g_{J_2+\epsilon_2/2, J_3}^{J+\epsilon_1/2} = U_{\epsilon_1, \epsilon_2}^{(J_3)}(J, J_2) g_{J_2 J_3}^J. \quad (2.34)$$

There are two types of flatness conditions. First, the inverse relations

$$U_{-\epsilon_1, -\epsilon_2}^{(J_3)}(J + \frac{\epsilon_1}{2}, J_2 + \frac{\epsilon_2}{2}) U_{\epsilon_1, \epsilon_2}^{(J_3)}(J, J_2) = 1.$$

Second, the elementary-plaquette relations

$$U_{\eta_1, \eta_2}^{(J_3)}(J + \frac{\epsilon_1}{2}, J_2 + \frac{\epsilon_2}{2}) U_{\epsilon_1, \epsilon_2}^{(J_3)}(J, J_2) = U_{\epsilon_1, \epsilon_2}^{(J_3)}(J + \frac{\eta_1}{2}, J_2 + \frac{\eta_2}{2}) U_{\eta_1, \eta_2}^{(J_3)}(J, J_2).$$

Each of the two types is enough to determine $g_{1/2 J}^{J-1/2}$, and we shall only detail the first one for $\epsilon_1 = \epsilon_2 = 1$. One has

$$U_{1,1}^{(J_3)}(J, J_2) = A \left[\begin{matrix} 1/2 & J_2 & J_2+1/2 \\ J_3 & J+1/2 & J \end{matrix} \right], \quad (2.35)$$

$$U_{-1,-1}^{(J_3)}(J + 1/2, J_2 + 1/2) = \frac{g_{1/2, J+1/2}^J}{g_{1/2, J_2+1/2}^{J_2}} A \left[\begin{matrix} 1/2 & J_2+1/2 & J_2 \\ J_3 & J & J+1/2 \end{matrix} \right]. \quad (2.36)$$

The inverse relation gives

$$g_{1/2 J_2+1/2}^{J_2} / g_{1/2 J+1/2}^J = A \left[\begin{matrix} 1/2 & J_2 & J_2+1/2 \\ J_3 & J+1/2 & J \end{matrix} \right] A \left[\begin{matrix} 1/2 & J_2+1/2 & J_2 \\ J_3 & J & J+1/2 \end{matrix} \right]. \quad (2.37)$$

So, the r.h.s. must be of a very specific form: it must be the quotient of a function of J by the same function of J_2 . Thanks to the particular form of the A coefficients, this is true, and one finds by explicit calculations that

$$g_{1/2 J_2+1/2}^{J_2} / g_{1/2 J+1/2}^J = \sqrt{\frac{F(1 + (2J_2 + 1)h/\pi)F(-1 - (2J_2 + 2)h/\pi)}{F(1 + (2J + 1)h/\pi)F(-1 - (2J + 2)h/\pi)}}.$$

The expression of $g_{1/2 J}^{J-1/2}$ is thereby determined up to an irrelevant constant factor (it does not affect F):

$$g_{1/2 J}^{J-1/2} = g_0 \sqrt{F(1 + 2Jh/\pi)F(-1 - (2J + 1)h/\pi)}. \quad (2.38)$$

It is straightforward to verify that the other flatness conditions are satisfied⁸. Finally, we solve the recurrence relations, and compute the general expression for $g_{J_1 J_2}^{J_{12}}$. Clearly, there exists an integer number p such that $(J_1 - p/2) + J_2 = J_{12} + p/2$, that is $p = J_1 + J_2 - J_{12}$. Thus, by repeated multiplications through $U_{-1,+1}^{(J_2)}$, we may relate $g_{J_1 J_2}^{J_{12}}$ to

$$g_{(J_1 - J_2 + J_{12})/2, J_2}^{(J_1 + J_2 + J_{12})/2} = 1.$$

This leads to

$$g_{J_1 J_2}^{J_{12}} = \prod_{n=1}^{J_1 + J_2 - J_{12}} U_{-1,+1}^{(J_2)}(J_{12} + n/2, J_1 - n/2).$$

Eqs.2.30, and 2.38 give

$$U_{-1,+1}^{(J_2)}(J_{12}, J_1) = g_0 \sqrt{\frac{F(1 + (2J_1 + 1)h/\pi)F(-1 - (2J_{12} + 1)h/\pi)}{F(1 + (J_1 + J_2 - J_{12})h/\pi)F((J_1 - J_2 - J_{12} - 1)h/\pi)}}.$$

After some re-arrangement, one finds

$$g_{J_1 J_2}^{J_{12}} = (g_0)^{J_1 + J_2 - J_{12}} \prod_{k=1}^{J_1 + J_2 - J_{12}} \sqrt{\frac{F(1 + (2J_1 - k + 1)h/\pi)F(1 + (2J_2 - k + 1)h/\pi)F(-1 - (2J_{12} + k + 1)h/\pi)}{F(1 + kh/\pi)}}. \quad (2.39)$$

The result is symmetric in J_1, J_2 . The lack of symmetry between J_1 or J_2 and J_{12} will be explained later on.

So far we only dealt with the fusing and braiding for $J_1 = 1/2$, and considered the matrix elements between highest-weight states. Next we show that the generalization to the full OPE follows from the associativity conditions, if we assume that Eq.2.22 holds for any matrix element, according to the MS scheme. Let us rederive, for pedagogical purpose, the associated pentagonal relation for the F matrices. We fuse $\langle \varpi_J | V_{m_1}^{(J_1)} V_{m_2}^{(J_2)} V_{m_3}^{(J_3)} | \varpi_{J+2m_1+2m_2+2m_3} \rangle$ in two different ways, beginning from the left, and from the right, and identify the coefficients of the resulting operator $V_{m_1+m_2+m_3}^{(J_{123}, \{\nu_{123}\})}$. This gives

$$\begin{aligned} & \sum_{J_{12}} F_{J+m_1+m_2, J_{123}} \left[\begin{matrix} J_{12} \\ J \end{matrix} \begin{matrix} J_3 \\ J+m_1+m_2+m_3 \end{matrix} \right] F_{J+m_1, J_{12}} \left[\begin{matrix} J_1 \\ J \end{matrix} \begin{matrix} J_2 \\ J+m_1+m_2 \end{matrix} \right] \times \\ & \sum_{\{\nu_{12}\}} \langle \varpi_{J_{123}}, \{\nu_{123}\} | V_{J_3-J_{123}}^{J_{12}, \{\nu_{12}\}} | \varpi_{J_3} \rangle \langle \varpi_{J_{12}}, \{\nu_{12}\} | V_{J_2-J_{12}}^{J_1} | \varpi_{J_2} \rangle = \\ & \sum_{J_{23}} F_{J+m_1, J_{23}} \left[\begin{matrix} J_1 \\ J \end{matrix} \begin{matrix} J_{23} \\ J+m_1+m_2+m_3 \end{matrix} \right] F_{J+m_1+m_2, J_{23}} \left[\begin{matrix} J_2 \\ J+m_1 \end{matrix} \begin{matrix} J_3 \\ J+m_1+m_2+m_3 \end{matrix} \right] \times \\ & \sum_{\{\nu_{23}\}} \langle \varpi_{J_{123}}, \{\nu_{123}\} | V_{J_{23}-J_{123}}^{J_1} | \varpi_{J_{23}}, \{\nu_{23}\} \rangle \langle \varpi_{J_{23}}, \{\nu_{23}\} | V_{J_3-J_{23}}^{J_2} | \varpi_{J_3} \rangle. \quad (2.40) \end{aligned}$$

On the r.h.s. we use

$$\sum_{\{\nu_{23}\}} | \varpi_{J_{23}}, \{\nu_{23}\} \rangle \langle \varpi_{J_{23}}, \{\nu_{23}\} | = \mathcal{P}_{J_{23}}, \quad (2.41)$$

⁸This only makes use of the condition $F(z)F(1-z) = 1$, and this construction is more general than the particular structure we are discussing.

and obtain the factor $\langle \varpi_{J_{123}}, \{\nu_{123}\} | V_{J_{23}-J_{123}}^{J_1} V_{J_3-J_{23}}^{J_2} | \varpi_{J_3} \rangle$. These last operators are fused in their turn, obtaining,

$$\begin{aligned} \sum_{J_{23}} F_{J+m_1+m_2, J_{23}} \left[\begin{matrix} J_2 & J_3 \\ J+m_1 & J+m_1+m_2+m_3 \end{matrix} \right] F_{J+m_1, J_{123}} \left[\begin{matrix} J_1 & J_{23} \\ J & J+m_1+m_2+m_3 \end{matrix} \right] F_{J_{23}, J_{12}} \left[\begin{matrix} J_1 & J_2 \\ J_{123} & J_3 \end{matrix} \right] \\ = F_{J+m_1, J_{12}} \left[\begin{matrix} J_1 & J_2 \\ J & J+m_1+m_2 \end{matrix} \right] F_{J+m_1+m_2, J_{123}} \left[\begin{matrix} J_{12} & J_3 \\ J & J+m_1+m_2+m_3 \end{matrix} \right]. \end{aligned} \quad (2.42)$$

On the other hand, the q-6j symbols are well known to satisfy the same relation[25], that is

$$\begin{aligned} \sum_{J_{23}} \left\{ \begin{matrix} J_2 & J_3 \\ J+m_1+m_2+m_3 & J+m_1 \end{matrix} \middle| \begin{matrix} J_{23} \\ J+m_1+m_2 \end{matrix} \right\} \left\{ \begin{matrix} J_1 & J_{23} \\ J+m_1+m_2+m_3 & J \end{matrix} \middle| \begin{matrix} J_{123} \\ J+m_1 \end{matrix} \right\} \left\{ \begin{matrix} J_1 & J_2 \\ J_3 & J_{123} \end{matrix} \middle| \begin{matrix} J_{12} \\ J_{23} \end{matrix} \right\} \\ = \left\{ \begin{matrix} J_1 & J_2 \\ J+m_1+m_2 & J \end{matrix} \middle| \begin{matrix} J_{12} \\ J+m_1 \end{matrix} \right\} \left\{ \begin{matrix} J_{12} & J_3 \\ J+m_1+m_2+m_3 & J \end{matrix} \middle| \begin{matrix} J_{123} \\ J+m_1+m_2 \end{matrix} \right\}. \end{aligned} \quad (2.43)$$

These two relations are used for a proof of Eq.2.28, by recurrence on J_1 , starting from our previous derivation of the case $J_1 = 1/2$. Assume that Eq.2.28 holds for $J_1 \leq K$, and write Eqs.2.42, and 2.43 for $J_1 \leq K$ and $J_2 \leq K$. In Eq.2.42, the last term, that is $F_{J+m_1+m_2, J_{123}} \left[\begin{matrix} J_{12} & J_3 \\ J & J+m_1+m_2+m_3 \end{matrix} \right]$ may have $J_{12} > K$, and then it is the only one which is not known from our hypothesis. Combining this equality with the similar one for 6-j symbols one immediately sees that $F_{J+m_1+m_2, J_{123}} \left[\begin{matrix} J_{12} & J_3 \\ J & J+m_1+m_2+m_3 \end{matrix} \right]$ is also given by Eq.2.28. Thus this relation holds for all J 's. $\square$

It is simple to see, at once that all the MS conditions will be satisfied since we may define new chiral vertex operators $\tilde{V}$, by

$$\mathcal{P}_{J_{12}} \tilde{V}_{J_2-J_{12}}^{J_1} \equiv g_{J_1 J_2}^{J_{12}} \mathcal{P}_{J_{12}} V_{J_2-J_{12}}^{J_1}, \quad (2.44)$$

so that the fusing and braiding matrices simply become

$$\tilde{F}_{J_{23}, J_{12}} \left[\begin{matrix} J_1 & J_2 \\ J & J_3 \end{matrix} \right] = \left\{ \begin{matrix} J_1 & J_2 \\ J_3 & J \end{matrix} \middle| \begin{matrix} J_{12} \\ J_{23} \end{matrix} \right\}, \quad \tilde{B}_{J_{23}, J_{12}}^\pm \left[\begin{matrix} J_1 & J_3 \\ J & J_2 \end{matrix} \right] = e^{\pm i\pi(\Delta_J + \Delta_{J_2} - \Delta_{J_{23}} - \Delta_{J_{12}})} \left\{ \begin{matrix} J_1 & J_2 \\ J_3 & J \end{matrix} \middle| \begin{matrix} J_{12} \\ J_{23} \end{matrix} \right\}. \quad (2.45)$$

It is well known[25] that the q-6j symbols satisfy the polynomial equations. For instance, we already used the pentagonal relation Eq.2.43. Let us stress, however, as already mentioned in the introduction, that **this does not mean that our fusing and braiding matrices are equivalent to Eqs.2.45 from the viewpoint of conformal theory**. Indeed, the new fusing equations read

$$\begin{aligned} \mathcal{P}_J \tilde{V}_{m_1}^{(J_1)} \tilde{V}_{m_2}^{(J_2)} = \mathcal{P}_J \sum_{J_{12}=|J_1-J_2|}^{J_1+J_2} \tilde{F}_{J+m_1, J_{12}} \left[\begin{matrix} J_1 & J_2 \\ J & J+m_1+m_2 \end{matrix} \right] \times \\ \sum_{\{\nu_{12}\}} \tilde{V}_{m_1+m_2}^{(J_{12}, \{\nu_{12}\})} \langle \varpi_{J_{12}}, \{\nu_{12}\} | \tilde{V}_{J_2-J_{12}}^{(J_1)} | \varpi_{J_2} \rangle. \end{aligned} \quad (2.46)$$

The matrix element of $\tilde{V}$ on the right-hand side is a book-keeping device to recover the coefficients of the OPE. There the normalization of $\tilde{V}$ appears explicitly. According to Eqs.2.44, Eq.2.11 is replaced by

$$\langle \varpi_{J_2} | \tilde{V}_m^{(J_1)}(1) | \varpi_{J_{12}} \rangle = g_{J_1 J_2}^{J_{12}} \delta_{m, J_{12}-J_2}. \quad (2.47)$$

For instance, at the level of primaries we have

$$\mathcal{P}_J \tilde{V}_{m_1}^{(J_1)} \tilde{V}_{m_2}^{(J_2)} = \mathcal{P}_J \sum_{J_{12}=|J_1-J_2|}^{J_1+J_2} g_{J_1 J_2}^{J_{12}} \tilde{F}_{J+m_1, J_{12}} \left[\begin{matrix} J_1 & J_2 \\ J & J+m_1+m_2 \end{matrix} \right] \tilde{V}_{m_1+m_2}^{(J_{12})} + \dots \quad (2.48)$$

The g 's have re-appeared. There is no way to get rid of them at the level of the two dimensional OPA.

Of course, it is easy to verify that the g factors of Eq.2.28 cancel in pairs, in every polynomial equation. Consider, for instance, the Yang-Baxter equation:

$$\begin{aligned} & \sum_{J_{134}} B_{J_{234}, J_{134}} \begin{bmatrix} J_1 & J_2 \\ J_{1234} & J_{34} \end{bmatrix} B_{J_{34}, J_{14}} \begin{bmatrix} J_1 & J_3 \\ J_{134} & J_4 \end{bmatrix} B_{J_{134}, J_{124}} \begin{bmatrix} J_2 & J_3 \\ J_{1234} & J_{14} \end{bmatrix} \\ &= \sum_{J_{24}} B_{J_{34}, J_{24}} \begin{bmatrix} J_2 & J_3 \\ J_{234} & J_4 \end{bmatrix} B_{J_{234}, J_{124}} \begin{bmatrix} J_1 & J_3 \\ J_{1234} & J_{24} \end{bmatrix} B_{J_{24}, J_{14}} \begin{bmatrix} J_1 & J_2 \\ J_{124} & J_4 \end{bmatrix}, \end{aligned} \quad (2.49)$$

which is obtained from operator-braidings in $\langle \varpi_{J_{1234}} | V^{(J_1)} V^{(J_2)} V^{(J_3)} | \varpi_{J_4} \rangle$. After cancelling out the g -factors, there remains

$$\begin{aligned} & \sum_{J_{134}} e^{i\pi\epsilon(\Delta_{J_{1234}} - \Delta_{J_{124}} - \Delta_{J_{134}} - \Delta_{J_{234}})} \left\{ \begin{matrix} J_1 & J_{34} \\ J_2 & J_{1234} \end{matrix} \middle| \begin{matrix} J_{134} \\ J_{234} \end{matrix} \right\} \left\{ \begin{matrix} J_1 & J_4 \\ J_3 & J_{134} \end{matrix} \middle| \begin{matrix} J_{14} \\ J_{34} \end{matrix} \right\} \left\{ \begin{matrix} J_2 & J_{14} \\ J_3 & J_{1234} \end{matrix} \middle| \begin{matrix} J_{124} \\ J_{134} \end{matrix} \right\} \\ &= \sum_{J_{24}} e^{i\pi\epsilon(\Delta_{J_4} - \Delta_{J_{14}} - \Delta_{J_{24}} - \Delta_{J_{34}})} \left\{ \begin{matrix} J_2 & J_4 \\ J_3 & J_{234} \end{matrix} \middle| \begin{matrix} J_{24} \\ J_{34} \end{matrix} \right\} \left\{ \begin{matrix} J_1 & J_{24} \\ J_3 & J_{1234} \end{matrix} \middle| \begin{matrix} J_{124} \\ J_{234} \end{matrix} \right\} \left\{ \begin{matrix} J_1 & J_4 \\ J_2 & J_{124} \end{matrix} \middle| \begin{matrix} J_{14} \\ J_{24} \end{matrix} \right\}, \end{aligned} \quad (2.50)$$

which is a relation satisfied by the 6-j coefficients [25].

As a conclusion of this section, we give the final form of the operator-algebra for the V fields. In the MS form, and according to Eqs.2.24 and 2.28, the fusing algebra reads

$$\begin{aligned} & \langle \varpi_{J_{123}}, \{\nu_{123}\} | V_{J_{23}-J_{123}}^{(J_1)} V_{J_3-J_{23}}^{(J_2)} | \varpi_{J_3}, \{\nu_3\} \rangle = \sum_{J_{12}=|J_1-J_2|}^{J_1+J_2} \frac{g_{J_1 J_2}^{J_{12}} g_{J_{12} J_3}^{J_{123}}}{g_{J_2 J_3}^{J_{23}} g_{J_1 J_{23}}^{J_{123}}} \left\{ \begin{matrix} J_1 & J_2 \\ J_3 & J_{123} \end{matrix} \middle| \begin{matrix} J_{12} \\ J_{23} \end{matrix} \right\} \times \\ & \sum_{\{\nu_{12}\}} \langle \varpi_{J_{123}}, \{\nu_{123}\} | V_{J_3-J_{123}}^{(J_{12}, \{\nu_{12}\})} | \varpi_{J_3}, \{\nu_3\} \rangle \langle \varpi_{J_{12}}, \{\nu_{12}\} | V_{J_2-J_{12}}^{(J_1)} | \varpi_{J_2} \rangle. \end{aligned} \quad (2.51)$$

In the operator formalism of the Liouville theory, this may be rewritten as

$$\begin{aligned} V_{m_1}^{(J_1)} V_{m_2}^{(J_2)} &= \sum_{J_{12}=|J_1-J_2|}^{J_1+J_2} \frac{g_{J_1 J_2}^{J_{12}} g_{J_{12}}^{(\varpi-\varpi_0)/2}}{g_{J_2}^{(\varpi-\varpi_0+2m_1+2m_2)/2} g_{J_1}^{(\varpi-\varpi_0)/2}} \times \\ & \left\{ \begin{matrix} J_1 & J_2 \\ (\varpi-\varpi_0+2m_1+2m_2)/2 & (\varpi-\varpi_0)/2 \end{matrix} \middle| \begin{matrix} J_{12} \\ (\varpi-\varpi_0+2m_1)/2 \end{matrix} \right\} \times \\ & \sum_{\{\nu_{12}\}} V_{m_1+m_2}^{(J_{12}, \{\nu_{12}\})} \langle \varpi_{J_{12}}, \{\nu_{12}\} | V_{J_2-J_{12}}^{(J_1)} | \varpi_{J_2} \rangle. \end{aligned} \quad (2.52)$$

In this last formula, ϖ is an operator. It is easy to check that this operator-expression is equivalent to Eq.2.51, by computing the matrix element between the states $\langle \varpi_{J_{123}}, \{\nu_{123}\} |$, and $| \varpi_{J_3}, \{\nu_3\} \rangle$. Then, the additional spins of Eq.2.51, as compared with Eq.2.52, are given by

$$\begin{aligned} J_{123} &= (\varpi - \varpi_0)/2, \\ J_{23} &= (\varpi - \varpi_0 + 2m_1)/2, \\ J_3 &= (\varpi - \varpi_0 + 2m_1 + 2m_2)/2. \end{aligned} \quad (2.53)$$

The explicit formula for the Racah-Wigner 6-j coefficients, which have the tetrahedral symmetry, are given in [25] by

$$\left\{ \begin{matrix} a & b & e \\ d & c & f \end{matrix} \right\} = ([z - a - b - e]! [z - a - c - f]!)^{-1/2} (-1)^{a+b-c-d-2e} \left\{ \begin{matrix} a & b \\ d & c \end{matrix} \middle| \begin{matrix} e \\ f \end{matrix} \right\} =$$

$$\begin{aligned}
&= \Delta(a, b, e) \Delta(a, c, f) \Delta(c, e, d) \Delta(d, b, f) \times \\
&\sum_{z \text{ integer}} (-1)^z [z+1]! \left[[z-a-b-e]! [z-a-c-f]! [z-b-d-f]! \times \right. \\
&\left. [z-d-c-e]! [a+b+c+d-z]! [a+d+e+f-z]! [b+c+e+f-z]! \right]^{-1} \quad (2.54)
\end{aligned}$$

with

$$\Delta(l, j, k) = \sqrt{\frac{[-l+j+k]! [l-j+k]! [l+j-k]!}{[l+j+k+1]!}}$$

and

$$[n]! \equiv \prod_{r=1}^n [r] \quad [r] \equiv \frac{\sin(hr)}{\sin h}. \quad (2.55)$$

For completeness, we also summarize the formulae for the braiding⁹. In the MS form, Eqs.2.25, and 2.26 show that it is given by

$$\begin{aligned}
&< \varpi_{J_{123}}, \{\nu_{123}\} | V_{J_{23}-J_{123}}^{(J_1)} V_{J_3-J_{23}}^{(J_2)} | \varpi_{J_3}, \{\nu_3\} > = \sum_{J_{13}} e^{\pm i\pi(\Delta_{J_{123}} + \Delta_{J_3} - \Delta_{J_{23}} - \Delta_{J_{13}})} \times \\
&\frac{g_{J_1 J_3}^{J_{13}} g_{J_{13} J_2}^{J_{123}}}{g_{J_2 J_3}^{J_{23}} g_{J_1 J_{23}}^{J_{123}}} \left\{ \begin{matrix} J_1 & J_3 \\ J_2 & J_{123} \end{matrix} \middle| \begin{matrix} J_{13} \end{matrix} \right\} < \varpi_{J_{123}}, \{\nu_{123}\} | V_{J_{13}-J_{123}}^{(J_2)} V_{J_3-J_{13}}^{(J_1)} | \varpi_{J_3}, \{\nu_3\} > . \quad (2.56)
\end{aligned}$$

The equivalent operator-form is given by

$$\begin{aligned}
V_{m_1}^{(J_1)} V_{m_2}^{(J_2)} &= \sum_{n_1+n_2=m_1+m_2} e^{\mp i h (2m_1 m_2 + m_2^2 - n_2^2 + \varpi(m_2 - n_2))} \times \\
&\left\{ \begin{matrix} J_1 & (\varpi - \varpi_0 + 2m_1 + 2m_2)/2 \\ J_2 & (\varpi - \varpi_0)/2 \end{matrix} \middle| \begin{matrix} (\varpi - \varpi_0 + 2n_2)/2 \\ (\varpi - \varpi_0 + 2m_1)/2 \end{matrix} \right\} \times \\
&\frac{g_{J_1}^{(\varpi - \varpi_0 + 2n_2)/2} g_{J_2}^{(\varpi - \varpi_0)/2}}{g_{J_1}^{(\varpi - \varpi_0 + 2n_1 + 2n_2)/2} g_{J_2}^{(\varpi - \varpi_0 + 2n_2)/2}} V_{n_2}^{(J_2)} V_{n_1}^{(J_1)}. \quad (2.57)
\end{aligned}$$

The correspondence table is again given by Eq.2.53, with, in addition,

$$J_{13} = (\varpi - \varpi_0 + 2n_2)/2. \quad (2.58)$$

In the operator-forms Eqs.2.52, 2.57, one sees that the fusion and braiding matrices involve the operator ϖ , and thus do not commute with the V-operators (see Eq.A.12). Such is the general situation of the operator-algebras in the MS formalism. This is in contrast with, for instance, the braiding relations for quantum group representations. In the coming section, and completing the results of refs.[13, 19], we will change basis to the holomorphic operators ξ which are such that these ϖ dependences of the fusing and braiding matrices disappear. After the transformation, one is in the same situation as for quantum group, and its structure becomes more transparent.

Our last point of the section will concern the treatment of the end points in correlators. Following ref.[27], one has

$$\begin{aligned}
&\lim_{z \rightarrow 0} V_{m \hat{m}}^{(J \hat{J})}(z) | \varpi_0 > = \delta_{m+J,0} \delta_{\hat{m}+\hat{J},0} | \varpi_{J, \hat{J}} > \\
&\lim_{z \rightarrow \infty} z^{2\Delta_{J \hat{J}}} < -\varpi_0 | V_{m \hat{m}}^{(J \hat{J})}(z) = \delta_{m+J,0} \delta_{\hat{m}+\hat{J},0} < -\varpi_{J, \hat{J}} |, \quad (2.59)
\end{aligned}$$

⁹an equivalent form is given in ref[30], without connection to the 6-j.

where $\Delta_{J\hat{J}}$ is the conformal weight (see Eq.A.15) and $\varpi_{J,\hat{J}} = \varpi_0 + 2J + 2\hat{J}\pi/h$.

At this point we consider shortly the general fields $V_{m\hat{m}}^{(J\hat{J})}$, since, keeping the $V_m^{(J)}$ fields only would give vanishing two-point functions. General correlators, which are $sl(2, C)$ invariant, may be described by the operator matrix-elements

$$\langle -\varpi_0 | V_{m_n \hat{m}_n}^{(J_n \hat{J}_n)}(z_n) \cdots V_{m_2 \hat{m}_2}^{(J_2 \hat{J}_2)}(z_2) V_{m_1 \hat{m}_1}^{(J_1 \hat{J}_1)}(z_1) | \varpi_0 \rangle. \quad (2.60)$$

If we are on the sphere, where the points 0 and ∞ are equivalent to any other points, the limit $z_1 \rightarrow 0$ must be finite, and the limit $z_n \rightarrow \infty$ must also become finite after transforming $z_n \rightarrow 1/z_n$. Thus, according to Eqs.2.59, one should only consider correlators with

$$m_n + J_n = \hat{m}_n + \hat{J}_n = m_1 + J_1 = \hat{m}_1 + \hat{J}_1 = 0. \quad (2.61)$$

Then the question arises whether the operator algebra just summarized has a consistent restriction, namely, whether, conditions Eq.2.61 are preserved under fusing and braiding. Of course, this is only possible since the F and B matrices are functions of ϖ . We shall only discuss the conditions arising from the limit $z_1 \rightarrow 0$ of the right-most operator, for the fields with $J_1 + m_1 = \hat{J}_1 + \hat{m}_1 = 0$, explicitly.

Consider first the fusion. One applies Eq.2.51, or equivalently, Eq.2.52, for the highest-weight state with $\varpi_{J_3} = \varpi_0$, so that $J_3 = 0$. Recall the defining relation of the 6-j symbols:

$$\sum_{J_{23}} (J_2, \mu_2; J_3, \mu_3 | J_{23}) (J_1, \mu_1; J_{23}, \mu_{23} | J_{123}) \left\{ \begin{matrix} J_1 & J_2 \\ J_3 & J_{123} \end{matrix} \middle| \begin{matrix} J_{12} \\ J_{23} \end{matrix} \right\} = \\ (J_1, \mu_1; J_2, \mu_2 | J_{12}) (J_{12}, \mu_{12}; J_3, \mu_3 | J_{123}). \quad (2.62)$$

The symbols $(J_k, \mu_k; J_\ell, \mu_\ell | J_{k\ell})$ denote the q Clebsch-Gordan (3-j) coefficients¹⁰. In our present situation, we have $J_3 = 0$, and thus $\mu_3 = 0$. Obviously one has $J_{123} = J_{12}$, and $J_{23} = J_3$, and one may easily verify, from the last relation, that

$$\left\{ \begin{matrix} J_1 & J_2 \\ 0 & J_{123} \end{matrix} \middle| \begin{matrix} J_{12} \\ J_{23} \end{matrix} \right\} = \delta_{J_{123}, J_{12}} \delta_{J_{23}, J_2}. \quad (2.63)$$

It follows from Eqs.2.53 that, for $J_3 = 0$, the fusing relation vanishes unless $m_2 + J_2 = 0$, that is, unless the restriction condition Eq.2.61 is satisfied. Then, the g -factor becomes one, and one gets

$$V_{m_1}^{(J_1)} V_{m_2}^{(J_2)} | \varpi_0 \rangle = \delta_{J_2+m_2, 0} \sum_{\{\nu_{12}\}} V_{m_1-J_2}^{(J_2-m_1, \{\nu_{12}\})} | \varpi_0 \rangle \langle \varpi_{J_2-m_2, \{\nu_{12}\}} | V_{J_2-J_{12}}^{(J_1)} | \varpi_{J_2} \rangle. \quad (2.64)$$

A similar discussion shows that

$$V_{m_1}^{(J_1)} V_{m_2}^{(J_2)} | \varpi_0 \rangle = \delta_{J_2+m_2, 0} e^{\mp i h [(\varpi_0 + 2J_2 - 2m_1)^2 + \varpi_0^2 - (\varpi_0 + 2J_2)^2 - (\varpi_0 + 2J_2)^2] / 4} \times \\ V_{m_1+J_1-J_2}^{(J_2)} V_{-J_1}^{(J_1)} | \varpi_0 \rangle. \quad (2.65)$$

Thus, the above restriction conditions are preserved by fusing and braiding. Note that one may equivalently describe the correlators by

$$\langle \varpi_0 | V_{m_n \hat{m}_n}^{(J_n \hat{J}_n)}(z_n) \cdots V_{m_2 \hat{m}_2}^{(J_2 \hat{J}_2)}(z_2) V_{m_1 \hat{m}_1}^{(J_1 \hat{J}_1)}(z_1) | -\varpi_0 \rangle.$$

Then, conditions Eq.2.61 are replaced by

$$m_n - J_n = \hat{m}_n - \hat{J}_n = m_1 - J_1 = \hat{m}_1 - \hat{J}_1 = 0.$$

The two formalisms are completely equivalent.

¹⁰We use a condensed notation $(J_1, \mu_1; J_2, \mu_2 | J_{12})$ instead of $(J_1, \mu_1; J_2, \mu_2 | J_{12}, \mu_1 + \mu_2)$.

3 THE COVARIANT OPERATOR ALGEBRA

In the preceding section, the quantum numbers J and m of the $V_m^{(J)}$ operators should be regarded as quantum-group invariant. Indeed, the J 's appear as total spins in q-6j symbols, and the m 's are given by differences of J 's, as is clear on Eq.2.11, for instance. Thus the quantum group does not act on the V 's. In refs.[13, 19], other fields $\xi_M^{(J)}$ were defined which are quantum-group covariant. Following ref.[19], this is done in two steps. A first change of field is performed by introducing ψ fields of the form

$$\psi_m^{(J)} \equiv E_m^{(J)}(\varpi) V_m^{(J)}, \quad (3.1)$$

such that the braiding matrix and the fusing coefficients for $\psi_\alpha^{(1/2)} \psi_m^{(J)} \rightarrow \psi_{m+\alpha}^{(J+1/2)}$ become trigonometric. The definition of $E_m^{(J)}(\varpi)$ (which in the notations of ref.[19] is equal to $C^{J-m, J+m} D^{J-m, J+m}(\varpi)$) is recalled in Eqs.A.16-A.25. Then the ξ fields are defined by expressions of the form

$$\xi_M^{(J)} = \sum_{-J \leq m \leq J} |J, \varpi\rangle_M^m \psi_m^{(J)}, \quad -J \leq M \leq J. \quad (3.2)$$

The explicit form of the coefficients $|J, \varpi\rangle_M^m$ is given in Eq.3.35, below. The formulae just written allow us to deduce the F and B matrices of the ξ fields from those of the V fields derived in the previous section. Indeed, it was already shown in ref[19] that the braiding matrix of the ξ field coincides with the universal R -matrix of $U_q(sl(2))$. Concerning the fusing matrices, we shall establish an explicit connection later on, by first relating the coefficients $|J, \varpi\rangle_M^m$ to a limit of q-Clebsch-Gordan coefficients. At the present stage of the discussion, it is more enlightening to proceed in another way. We shall first transform the fusing matrix for the OPE of $V_{m_1}^{(1/2)}$ and $V_{m_2}^{(J_2)}$, which was the starting point of last section, and after, generalize the result using the associativity of the OPA.

Consider, thus Eq.2.24, with $J_1 = 1/2$, and make use of Eq.2.27. Taking Eq.A.12 into account, one sees that it is appropriate to multiply both sides by $E_{\pm 1/2}^{(1/2)}(\varpi) E_m^{(J)}(\varpi \pm 1)$. Using the recurrence relations satisfied by C and D (Eqs.A.15, and A.17 of ref.[19]) one thereby derives the fusing relations

$$\begin{aligned} \psi_{\pm 1/2}^{(1/2)} \psi_m^{(J)} &= \sum_{\{\nu\}} \frac{[\varpi \mp J + m]}{[\varpi]} \psi_{m \pm 1/2}^{(J+1/2, \{\nu\})} < \varpi_J + 1, \{\nu\} | V_{-1/2}^{(1/2)} | \varpi_J > . \\ &+ \sum_{\{\nu\}} \frac{[\mp J + m]}{[\varpi]} \Gamma(1 + 2Jh/\pi) \Gamma(-1 - (2J+1)h/\pi) \times \\ &\psi_{m \pm 1/2}^{(J-1/2, \{\nu\})} < \varpi_J - 1, \{\nu\} | V_{1/2}^{(1/2)} | \varpi_J > \end{aligned} \quad (3.3)$$

The original motivation for introducing the ψ fields[19, 20] was that the braiding matrix, and the leading-order fusing coefficients for them are trigonometrical. The last equation written shows that this is not true for the other fusing coefficients $\psi_\alpha^{(1/2)} \psi_m^{(J)} \rightarrow \psi_{m+\alpha}^{(J-1/2)}$. As a result, the OPE is not associative if one forgets the contribution of the secondaries (more about this below). Next, comparing with the expression Eq.2.38 of the g 's, one sees that the last equation is naturally rewritten as

$$\psi_{\pm 1/2}^{(1/2)} \psi_m^{(J)} = \sum_{\epsilon, \{\nu\}} g_{1/2, J}^{J+\epsilon/2} N_{\pm 1/2, m; m \pm 1/2}^{1/2, J; J+\epsilon/2; \varpi} \psi_{m \pm 1/2}^{(J+\epsilon/2, \{\nu\})} < \varpi_J + \epsilon, \{\nu\} | V_{-\epsilon/2}^{(1/2)} | \varpi_J > \quad (3.4)$$

where,

$$N_{\pm 1/2, m; m \pm 1/2}^{1/2, J; J+1/2; \varpi} = \frac{[\varpi \mp J + m]}{[\varpi]},$$

$$N_{\pm 1/2, m; m \pm 1/2}^{1/2, J; J-1/2; \varpi} = \frac{i\pi g_0^{-1}}{\sqrt{\sin(2hJ) \sin((2J+1)h)}} \frac{[\mp J + m]}{[\varpi]}. \quad (3.5)$$

One sees that the non-trigonometric part is entirely contained in the g 's. It will be shown at the end of the section that Eq.3.4 is a particular case of the general fusing algebra

$$\psi_{m_1}^{(J_1)} \psi_{m_2}^{(J_2)} = \sum_{J, \{\nu\}} g_{J_1, J_2}^J N_{m_1, m_2; m_1+m_2}^{J_1, J_2; J} \varpi \left| \psi_{m_1+m_2}^{(J, \{\nu\})} < \varpi_J, \{\nu\} | V_{J_2-J}^{(J_1)} | \varpi_{J_2} > . \quad (3.6)$$

In addition, $N_{m_1, m_2; m_1+m_2}^{J_1, J_2; J} \varpi$ will be related to a q-6j symbol. For the time being we transform Eq.3.4 further, in order to derive the fusion of the ξ fields. Their definition Eq.3.2, together with the shift properties of the V fields (see Eq.2.11, or Eq.A.12), are such that

$$\xi_\alpha^{(1/2)} \xi_M^{(J)} = \sum_{\eta=\pm 1/2, m} |1/2, \varpi)_\alpha^\eta | J, \varpi + 2\eta)_M^m \psi_\eta^{(1/2)} \psi_m^{(J)}. \quad (3.7)$$

It will be shown in appendix C that, if we choose $g_0 = 2\pi$, we have

$$\sum_{\eta=\pm 1/2, m, (\eta+m=n)} |1/2, \varpi)_\alpha^\eta | J, \varpi + 2\eta)_M^m N_{\eta, m; \eta+m}^{1/2, J; J+\epsilon/2; \varpi}$$

$$= (1/2, \alpha; J, M | J + \epsilon/2) | J + \epsilon/2, \varpi)_{M+\alpha}^n. \quad (3.8)$$

(recall that $(1/2, \alpha; J, M | J + \epsilon/2)$ denotes the 3-j symbols). Eq.3.4 becomes

$$\xi_{\pm 1/2}^{(1/2)} \xi_m^{(J)} = \sum_{\epsilon=\pm 1} g_{1/2, J}^{J+\epsilon/2} (1/2, \alpha; J, M | J + \epsilon/2) \times$$

$$\sum_{\{\nu\}} \xi_{m \pm 1/2}^{(J+\epsilon/2, \{\nu\})} < \varpi_J + \epsilon/2, \{\nu\} | V_{-\epsilon/2}^{(1/2)} | \varpi_J > . \quad (3.9)$$

Our next task is to generalize this last fusing identity. Note an important feature of this equation. As expected, it expresses the OPE of two ξ fields in terms of one ξ fields, (and its descendants), that is $\xi_{m \pm 1/2}^{(J+\epsilon/2, \{\nu\})}$. However, the coefficients of this OPE are proportional to the matrix element of a V field. The basic reason is that the V matrix element, of the fusing equation for the V fields (Eq.2.24) contains no m_1 or m_2 dependence, and is thus unchanged when going from the ψ to the ξ fields. Thus we shall start from the general ansatz

$$\mathcal{P}_J \xi_{M_1}^{(J_1)} \xi_{M_2}^{(J_2)} = \mathcal{P}_J \sum_{J_{12}=|J_1-J_2|}^{J_1+J_2} \mathcal{F}(J_1, M_1, J_2, M_2, J_{12}, M_{12}, \varpi_J) \times$$

$$\sum_{\{\nu\}} \xi_{M_1+M_2}^{(J_{12}, \{\nu\})} < \varpi_{J_{12}}, \{\nu\} | V_{J_2-J_{12}}^{(J_1)} | \varpi_{J_2} > . \quad (3.10)$$

Our next task is to prove that the fusion coefficients are proportional to the CG coefficients and independent of ϖ_J :

$$\mathcal{F}(J_1, M_1, J_2, M_2, J_{12}, M_{12}, \varpi_J) = g_{J_1 J_2}^{J_{12}}(J_1, M_1; J_2, M_2 | J_{12}). \quad (3.11)$$

To do this, we have to write the associativity equations for the ξ fields. The demonstration is the same as the one for the V operators (Eq.2.42), this is why we will not give all the details right now (more about it soon, however). There is a difference yet, that we have to emphasize. In the demonstration of Eq.2.42, we pointed out that only four of the five fusing coefficients really came from the fusion of the operators considered, as the other one — the third one on the left-hand side of Eq.2.42 — came from the fusion of the operators in the matrix elements, which had been restored as operators thanks to the closure relation. So, in the case of the ξ operators, the four fusion coefficients will become coefficients of the ξ , but **the other one will remain a fusing coefficient of V-fields**. Accordingly, we get a pentagonal relation of the form

$$\sum \mathcal{F} \mathcal{F} F = \mathcal{F} \mathcal{F}. \quad (3.12)$$

Next, using the explicit expression Eq.3.10, let us show that Eq.3.11 is a solution of the associativity condition. Indeed, with this ansatz, the pentagonal relation becomes

$$\begin{aligned} \sum_{J_{23}} (J_2, M_2; J_3, M_3 | J_{23}) (J_1, M_1; J_{23}, M_{23} | J_{123}) \left\{ \begin{matrix} J_1 & J_2 \\ J_3 & J_{123} \end{matrix} \middle| \begin{matrix} J_{12} \\ J_{23} \end{matrix} \right\} = \\ (J_1, M_1; J_2, M_2 | J_{12}) (J_{12}, M_{12}; J_3, M_3 | J_{123}). \end{aligned} \quad (3.13)$$

This is the basic identity that defines the q-6j coefficients[25], as already recalled (see Eq.2.62). Once we know that the equation 3.11 gives an associative algebra, it is easy to derive it by recursion from the particular case $J_1 = 1/2$, J_2 arbitrary (Eq.3.9) using a recurrence proof which is completely parallel to the one we gave for the V fields, in the previous section. We do not go through it again, and consider Eq.3.11 as established. $\square$

An important feature of the result is that the fusing matrix does not depend upon ϖ_J . Thus the projector of Eq.3.10 does not serve any purpose and may be removed. The fusion algebra finally reads

$$\begin{aligned} \xi_{M_1}^{(J_1)} \xi_{M_2}^{(J_2)} = \sum_{J_{12}=|J_1-J_2|}^{J_1+J_2} g_{J_1 J_2}^{J_{12}} (J_1, M_1; J_2, M_2 | J_{12}) \times \\ \sum_{\{\nu\}} \xi_{M_1+M_2}^{(J_{12}, \{\nu\})} < \varpi_{J_{12}}, \{\nu\} | V_{J_2-J_{12}}^{(J_1)} | \varpi_{J_2} >. \end{aligned} \quad (3.14)$$

For the coming discussion, we shall actually need the following generalization of this last relation

$$\begin{aligned} \xi_{M_1}^{(J_1, \{\gamma\})} \xi_{M_2}^{(J_2, \{\mu\})} = \sum_{J_{12}=|J_1-J_2|}^{J_1+J_2} g_{J_1 J_2}^{J_{12}} (J_1, M_1; J_2, M_2 | J_{12}) \times \\ \sum_{\{\nu\}} \xi_{M_1+M_2}^{(J_{12}, \{\nu\})} < \varpi_{J_{12}}, \{\nu\} | V_{J_2-J_{12}}^{(J_1, \{\gamma\})} | \varpi_{J_2}, \{\mu\} >, \end{aligned} \quad (3.15)$$

which holds according to the general principles of ref.[10].

We have just shown that the complete fusion rule of two ξ fields is most naturally expressed in terms of one ξ field and one V field. Now, we use this fact to operatorially relate the fusing and braiding properties of V fields with those of

the ξ fields. This will provide the general identities which relate their fusing and braiding matrices. The method is to apply the fusion algebra Eq.3.15 repeatedly to the OPE of several ξ fields. In fact, the forthcoming calculation will explicitly verify some of the polynomial equations of the OPE of the ξ fields, and may also be regarded as a pedagogical explanation of the arguments given above to derive Eq.3.15. For that purpose, one should of course deal with the descendant matrix-elements, and this is why Eq.3.15 is needed. Consider the matrix element $\langle \varpi_J, \{\alpha\} | \xi_{M_1}^{(J_1)} \xi_{M_2}^{(J_2)} \xi_{M_3}^{(J_3, \{\rho\})} | \varpi_K, \{\beta\} \rangle$. We apply Eq.3.15 twice: $\xi_{M_3}^{(J_3, \{\rho\})}$ and $\xi_{M_2}^{(J_2)}$ are fused first, and the result is then fused with $\xi_{M_1}^{(J_1)}$. One gets

$$\begin{aligned} & \langle \varpi_J, \{\alpha\} | \xi_{M_1}^{(J_1)} \xi_{M_2}^{(J_2)} \xi_{M_3}^{(J_3, \{\rho\})} | \varpi_K, \{\beta\} \rangle = \\ & \sum_{J_{23}\{\nu\}} g_{J_2 J_3}^{J_{23}} (J_2, M_2; J_3, M_3 | J_{23}) \sum_{J_{123}\{\mu\}} g_{J_1 J_{23}}^{J_{123}} (J_1, M_1; J_{23}, M_2 + M_3 | J_{123}) \times \\ & \langle \varpi_J, \{\alpha\} | \xi_{M_1+M_2+M_3}^{(J_{123}, \{\mu\})} | \varpi_K, \{\beta\} \rangle \times \\ & \langle \varpi_{J_{123}}, \{\mu\} | V_{J_{23}-J_{123}}^{(J_1)} | \varpi_{J_{23}}, \{\nu\} \rangle \langle \varpi_{J_{23}}, \{\nu\} | V_{J_3-J_{23}}^{(J_2)} | \varpi_{J_3}, \{\rho\} \rangle. \end{aligned}$$

Performing the sum over $\{\nu\}$ gives

$$\begin{aligned} & \langle \varpi_J, \{\alpha\} | \xi_{M_1}^{(J_1)} \xi_{M_2}^{(J_2)} \xi_{M_3}^{(J_3, \{\rho\})} | \varpi_K, \{\beta\} \rangle = \\ & \sum_{J_{123} J_{23}\{\nu\}} g_{J_1 J_{23}}^{J_{123}} (J_1, M_1; J_{23}, M_2 + M_3 | J_{123}) g_{J_2 J_3}^{J_{23}} (J_2, M_2; J_3, M_3 | J_{23}) \times \\ & \langle \varpi_J, \{\alpha\} | \xi_{M_1+M_2+M_3}^{(J_{123}, \{\nu\})} | \varpi_K, \{\beta\} \rangle \langle \varpi_{J_{123}}, \{\nu\} | V_{J_{23}-J_{123}}^{(J_1)} V_{J_3-J_{23}}^{(J_2)} | \varpi_{J_3}, \{\rho\} \rangle. \quad (3.16) \end{aligned}$$

At this point it is convenient to consider the last equation as follows. The two fields $\xi_{M_1}^{(J_1)}$ and $\xi_{M_2}^{(J_2)}$ on the left-hand side have been converted into two V fields, that is $V_{J_{23}-J_{123}}^{(J_1)}$, and $V_{J_3-J_{23}}^{(J_2)}$, on the right-hand side. The third field $\xi_{M_3}^{(J_3, \{\rho\})}$ plays the role of a background field which allows us to operatorially relate ξ fields to V fields by successive fusions. Its quantum numbers J_3 and $\{\rho\}$ specify which matrix element of V operators will come out at the end. Its quantum number M_3 is arbitrary and does not appear in the final matrix element of the two V fields. We shall come back to it later on. It is easily seen that this procedure may be repeated for more than three ξ fields, and that the structure is similar. The right-most ξ field which is the only one not converted into a V field is to be considered as a background field. In fact, each V field is multiplied by a 3-j symbols, and this method naturally leads to the transformation through dressing by 3-j symbols of refs.[12, 11], as we see next. Clearly, fusing or braiding the ξ fields and the corresponding V fields on each side of Eq.3.16 will directly relate their F and B matrices. Since this relation is, to begin with, different from the one which comes out from the connection through the $|J, \varpi\rangle_M^m$ coefficients, we next go through the derivations.

First consider the fusion. It follows from Eq.2.51 that the right member of Eq.3.16 may be rewritten as

$$\begin{aligned} & \sum_{J_{123} J_{23}} (J_1, M_1; J_{23}, M_2 + M_3 | J_{123}) (J_2, M_2; J_3, M_3 | J_{23}) \left\{ \begin{matrix} J_1 & J_2 & \\ J_3 & J_{123} & \end{matrix} \middle| \begin{matrix} J_{12} \\ J_{23} \end{matrix} \right\} g_{J_1 J_2}^{J_{12}} g_{J_{12} J_3}^{J_{123}} \times \\ & \sum_{\{\nu\}} \langle \varpi_J, \{\alpha\} | \xi_{M_1+M_2+M_3}^{(J_{123}, \{\nu\})} | \varpi_K, \{\beta\} \rangle \times \end{aligned}$$

$$\sum_{\{\gamma\}} \langle \varpi_{J_{123}}, \{\nu\} | V_{J_3-J_{123}}^{(J_{12}, \{\gamma\})} | \varpi_{J_3}, \{\rho\} \rangle \langle \varpi_{J_{12}}, \{\gamma\} | V_{J_2-J_{12}}^{(J_1)} | \varpi_{J_2} \rangle . \quad (3.17)$$

On the left-hand side, we fuse $\xi_{M_1}^{(J_1)}$, and $\xi_{M_2}^{(J_2)}$, making use of Eq.3.15. This gives

$$\sum_{J_{12}} g_{J_1 J_2}^{J_{12}}(J_1, M_1; J_2, M_2 | J_{12}) \sum_{\{\gamma\}} \langle \varpi_J, \{\alpha\} | \xi_{M_1+M_2}^{(J_{12}, \{\gamma\})} \xi_{M_3}^{(J_3, \{\rho\})} | \varpi_K, \{\beta\} \rangle \times \\ \langle \varpi_{J_{12}}, \{\gamma\} | V_{J_2-J_{12}}^{(J_1)} | \varpi_{J_2} \rangle . \quad (3.18)$$

Next the remaining two ξ fields are fused in their turn, and one gets

$$\sum_{J_{12}, J_{123}} g_{J_1 J_2}^{J_{12}}(J_1, M_1; J_2, M_2 | J_{12}) g_{J_{12} J_3}^{J_{123}}(J_{12}, M_1 + M_2; J_3, M_3 | J_{123}) \times \\ \sum_{\{\gamma\} \{\nu\}} \langle \varpi_J, \{\alpha\} | \xi_{M_1+M_2+M_3}^{(J_{123}, \{\nu\})} | \varpi_K, \{\beta\} \rangle \times \\ \langle \varpi_{J_{123}}, \{\nu\} | V_{J_3-J_{123}}^{(J_{12}, \{\gamma\})} | \varpi_{J_3}, \{\rho\} \rangle \langle \varpi_{J_{12}}, \{\gamma\} | V_{J_2-J_{12}}^{(J_1)} | \varpi_{J_2} \rangle . \quad (3.19)$$

Comparing this last expression with Eq.3.17, one sees that they coincide if the following relation holds:

$$\sum_{J_{23}} (J_2, M_2; J_3, M_3 | J_{23}) (J_1, M_1; J_{23}, M_2 + M_3 | J_{123}) \left\{ \begin{matrix} J_1 & J_2 \\ J_3 & J_{123} \end{matrix} \middle| \begin{matrix} J_{12} \\ J_{23} \end{matrix} \right\} = \\ (J_1, M_1; J_2, M_2 | J_{12}) (J_{12}, M_1 + M_2; J_3, M_3 | J_{123}). \quad (3.20)$$

This is indeed the defining relation of the 6-j symbols already recalled in Eq.2.62. Thus Eq.3.16 does establish the correct correspondence between the fusion properties of the V and ξ fields.

Consider, next, the braiding. For the V fields, it is given by Eq.2.56, or 2.57. Concerning the ξ fields, the braiding properties were derived in ref.[19]. One has

$$\xi_{M_1}^{(J_1)} \xi_{M_2}^{(J_2)} = \sum_{-J_1 \leq N_1 \leq J_1; -J_2 \leq N_2 \leq J_2} (J_1, J_2)_{M_1 M_2}^{N_2 N_1} \xi_{N_2}^{(J_2)} \xi_{N_1}^{(J_1)}. \quad (3.21)$$

The symbol $(J_1, J_2)_{M_1 M_2}^{N_2 N_1}$ denotes the following matrix element of the universal R-matrix:

$$(J_1, J_2)_{M_1 M_2}^{M'_2 M'_1} = \left(\ll J_1, M_1 | \otimes \ll J_2, M_2 | \right) \mathbf{R} \left(| J_1, M'_1 \gg \otimes | J_2, M'_2 \gg \right), \quad (3.22)$$

where $|J, M \gg$ are group theoretic states which span the representation of spin J of $U_q(sl(2))$. The universal R-matrix R is given by

$$\mathbf{R} = e^{-2ihJ_3 \otimes J_3} \sum_{n=0}^{\infty} \frac{(1 - e^{2ih})^n e^{ihn(n-1)/2}}{[n]!} e^{-ihnJ_3} (J_+)^n \otimes e^{ihnJ_3} (J_-)^n, \quad (3.23)$$

$J_{\pm}$, and J_3 are the quantum-group generators. For later use we recall that the R -matrix-elements may be simply written in terms of CG coefficients, since the latter are “twisted” eigenvectors, namely,

$$\sum_{N_1 N_2} (J_2, N_2; J_1, N_1 | J_{12}) (J_1, J_2)_{M_1 M_2}^{N_2 N_1} = e^{i\pi(\Delta_{J_{12}} - \Delta_{J_1} - \Delta_{J_2})} (J_1, M_1; J_2, M_2 | J_{12}). \quad (3.24)$$

It follows from the orthogonality of the CG coefficients that

$$(J_1, J_2)_{M_1 M_2}^{N_2 N_1} = \sum_{J_{12}} (J_2, N_2; J_1, N_1 | J_{12}) e^{i\pi(\Delta_{J_{12}} - \Delta_{J_1} - \Delta_{J_2})} (J_1, M_1; J_2, M_2 | J_{12}). \quad (3.25)$$

Returning to our main line, we follow the same procedure as for fusion. We shall skip details since the present discussion goes in close parallel. One exchanges the first two ξ fields on the left-hand side of Eq.3.16, and the two V fields on its right-hand side. Comparing the results, one derives the consistency condition

$$\begin{aligned} \sum_{N_1 N_2} (J_1, N_1; J_3, M_3 | J_{13}) (J_2, N_2; J_{13}, N_1 + M_3 | J_{123}, M_{123}) (J_1, J_2)_{M_1 M_2}^{N_2 N_1} = \\ \sum_{J_{23}} (J_2, M_2; J_3, M_3 | J_{23}) (J_1, M_1; J_{23}, M_2 + M_3 | J_{123}) \times \\ e^{i\pi(\Delta_{J_{123}} + \Delta_{J_3} - \Delta_{J_{23}} - \Delta_{J_{13}})} \left\{ \begin{matrix} J_1 & J_3 & J_{13} \\ J_2 & J_{123} & J_{23} \end{matrix} \right\}. \end{aligned} \quad (3.26)$$

This last relation may be easily proven using Eq.2.50 with $J_4 = 0$, Eq.2.62, and Eq.3.25. This defines the 6-j coefficient of the second type as was introduced in ref.[32].

The outcome of the present discussion is that the defining relations for the two types of 6-j symbols (Eqs.3.20 and 3.26) may be considered as relating the braiding and fusing matrices of the V and ξ fields. Thus the connection is established in a way where the quantum-group meaning is transparent. Clearly, Eqs.3.20, and 3.26 show that the connection is established via 3-j symbols. As a matter of fact, we have effectively re-derived the transformation through dressing by 3-j symbols of refs.[12, 11]. This is in contrast with the intrinsic transformation of refs.[13, 19], recalled in Eq.3.2, which uses the $|J, \varpi\rangle_M^m$ coefficients. Our next point is to establish the connection between these two transformations. The q-CG symbols involve 5 independent quantum numbers, and the $|J, \varpi\rangle_M^m$ coefficients only 4. In this connection, we have remarked that, in Eq.3.16, M_3 does not appear in the V -matrix element. It only appears in the two CG coefficients. The first one, that is $(J_1, M_1; J_{23}, M_2 + M_3 | J_{123})$, (resp. the second one, that is $(J_2, M_2; J_3, M_3 | J_{23})$) only involve J-quantum-numbers of the field $V_{J_{23}-J_{123}}^{(J_1)}$, (resp. $V_{J_3-J_{23}}^{(J_2)}$). These two sets of quantum numbers are treated on the same footing, but this is not the case for the M -quantum numbers, however, since the first CG coefficient contains $M_1, M_2 + M_3$, and the second M_2, M_3 . To motivate the coming mathematical derivation, we may remark that the symmetry is restored if $M_2 + M_3 \sim M_3$, that is if M_3 is very large compared with M_2 . One way to achieve this is to keep J_3 finite and to continue the quantum group for $M_3 > J_3$. This may be done rigourously, as we shall next show, since the q-3j symbols are given by q-deformed hypergeometric functions[25, 20]. In this limit one quantum number of the q-CG symbol drops out, and we will be left with the right number to identify the result with a $|J, \varpi\rangle_M^m$ coefficient.

We start from the explicit expression of the CG coefficients[25, 20], that is,

$$\begin{aligned} (J_1, M_1; J_2, M_2 | J) = B(J_1, J_2, J, M_1) \times \\ \sqrt{[J_2 - M_2]! [J_2 + M_2]! [J - M_1 - M_2]! [J + M_1 + M_2]!} \times \\ e^{ihM_2 J_1} \sum_{\mu=0}^{J_1+J_2-J} \left\{ \frac{e^{-ih\mu(J+J_1+J_2+1)} (-1)^\mu}{[\mu]! [J_1 + J_2 - J - \mu]!} \right\} \times \end{aligned}$$

$$\frac{1}{[J_1 - M_1 - \mu]! [J - J_2 + M_1 + \mu]! [J_2 + M_2 - \mu]! [J - J_1 - M_2 + \mu]!} \}, \quad (3.27)$$

where

$$B(J_1, J_2, J, M_1) = e^{ih(J_1+J_2-J)(J_1+J_2+J+1)/2} e^{-ihM_1J_2} \sqrt{[J_1 - M_1]! [J_1 + M_1]!} \times \\ \sqrt{[2J+1]} \sqrt{\frac{[J_1 + J_2 - J]! [-J_1 + J_2 + J]! [J_1 - J_2 + J]!}{[J_1 + J_2 + J + 1]!}}. \quad (3.28)$$

The terms in M_2 are conveniently rewritten as

$$\sqrt{\frac{[J_2 - M_2]!}{[J - J_1 + \mu - M_2]!}} \sqrt{\frac{[J - M_1 - M_2]!}{[J - J_1 + \mu - M_2]!}} \sqrt{\frac{[J_2 + M_2]!}{[J_2 - \mu + M_2]!}} \sqrt{\frac{[J + M_1 + M_2]!}{[J_2 - \mu + M_2]!}}. \quad (3.29)$$

We shall take the limit by giving an imaginary part to M_2 , thus we have to continue the above formulae in this variable. The last formula contains all the M_2 dependence. It has been written as a product of square roots of ratios of q-deformed factorials. Consider each term one-by-one. the differences between the arguments of numerators and denominators are

$$\begin{cases} J_2 - M_2 - (J - J_1 + \mu - M_2) & = J_1 + J_2 - J - \mu \\ J - M_1 - M_2 - (J - J_1 + \mu - M_2) & = J - M_1 - \mu \\ J_2 + M_2 - (J_2 - \mu + M_2) & = \mu \\ J + M_1 + M_2 - (J_2 - \mu + M_2) & = J + M_1 - J_2 + \mu. \end{cases}$$

The right members are independent from M_2 . In the expression Eq.3.27, the actual range of summation is dictated by the fact that a factorial with negative argument is infinite, so that each factorial may only have a non-negative argument. This immediately shows that the right-hand sides of the last set of equations are non-negative integers. As in ref.[20], let us introduce (ν is a positive integer, and a arbitrary)

$$[a]_\nu \equiv [a][a+1] \cdots [a+\nu-1] = \frac{[a+\nu-1]!}{[a-1]!}. \quad (3.30)$$

Eq.3.29 may be rewritten as

$$\sqrt{[J - J_1 + \mu - M_2 + 1]_{J_1+J_2-J-\mu} [J - J_1 + \mu - M_2 + 1]_{J-M_1-\mu}} \times \\ \sqrt{[J_2 - \mu + M_2 + 1]_\mu [J_2 - \mu + M_2 + 1]_{J+M_1-J_2+\mu}}. \quad (3.31)$$

According to the definition Eq.3.30, each term involves a number of factors which is independent from M_2 , so that the last expression makes sense for arbitrary complex M_2 . The limit is taken with an imaginary part, since, otherwise, the functions $\sin[h(M_2 + \alpha)]$, α constant, which appear in Eq.3.31, would not have a well defined limit. Of course, with the imaginary part, one exponential of trigonometric functions blows up while the other vanishes. The choice of sign is such that

$$\lim_{M_2 \rightarrow \infty} [\alpha \pm M_2] = \mp \frac{1}{2i \sin h} e^{-ih(M_2 \pm \alpha)} \quad (3.32)$$

$$\frac{[\alpha \pm M_2]!}{[\beta \pm M_2]!} \sim (-1)^{(\alpha-\beta)(1\pm 1)/2} (2i \sin h)^{\beta-\alpha} e^{ih(\beta-\alpha)M_2} e^{\mp ih(\alpha-\beta)(\alpha+\beta+1)/2}. \quad (3.33)$$

Substitute into Eq.3.27, one gets

$$(J_1, M_1; J_2, M_2|J) \sim e^{ih[-M_1(J+1/2)+(J_2^2-J_1^2-J^2+2J_1J_2)/2+(J_1+J_2-J)/2]} \times$$

$$B(J_1, J_2, J, M_1) \frac{e^{i\pi(J-J_2+M_1)/2}}{(2i \sin h)^{J_1}} \sum_{\mu=0}^{J_1+J_2-J} \left\{ \frac{e^{-2ih\mu(J+J_2+1)}}{[\mu]! [J_1+J_2-J-\mu]!} \times \right.$$

$$\left. \frac{1}{[J_1-M_1-\mu]! [J-J_2+M_1+\mu]!} \right\}. \quad (3.34)$$

Note that the limit is perfectly finite, since all exponentials in M_2 cancell out. On the other hand, the explicity expression of $|j, \varpi\rangle_M^m$ is [19]¹¹

$$|j, \varpi\rangle_M^m = \sqrt{\binom{2j}{j+M}} e^{ihm/2} \times$$

$$\sum_t \frac{e^{iht(\varpi+m)} [j-M]! [j+M]!}{\left[\frac{j-M+m-t}{2} \right]! \left[\frac{j-M-m+t}{2} \right]! \left[\frac{j+M+m+t}{2} \right]! \left[\frac{j+M-m-t}{2} \right]!} \quad (3.35)$$

and, letting $\mu = (j+M-m-t)/2$,

$$|j, \varpi\rangle_M^m = \sqrt{\binom{2j}{j+M}} e^{ihm/2} e^{ih((\varpi+m)(j+M-m))} \times$$

$$\sum_{\mu} \frac{e^{-2ih\mu(\varpi+m)} [j-M]! [j+M]!}{[\mu]! [j-m-\mu]! [m-M+\mu]! [j+M-\mu]!}. \quad (3.36)$$

Comparing Eqs.3.34, and 3.36, one sees that the variables should be related by

$$\varpi = \varpi_0 + 2J_2, \quad J_1 = j, \quad M = -M_1, \quad m = J - J_2. \quad (3.37)$$

One gets altogether,

$$(J_1, M_1; J_2, M_2|J) \sim \sqrt{\frac{[2J+1]}{\binom{2J_1}{J_1+J_2-J} [J_2+J-J_1+1]_{2J_1+1}}} \times$$

$$e^{i\pi[(J-J_2+M_1)/2+J+M_1-J_1-J_2]} \frac{e^{ih(M_1+J_2-J)/2}}{(2i \sin h)^{J_1}} |J_1, \varpi_0 + 2J_2\rangle_{-M_1}^{J-J_2}. \quad (3.38)$$

Recall that one has [20]

$$\sum_{M=-j}^j (-1)^{j-M} e^{ih(j-M)} |j, \varpi\rangle_M^m |j, \varpi + 2p\rangle_{-M}^{-p} = \delta_{m,p} C_m^{(j)}(\varpi), \quad (3.39)$$

where

$$C_m^{(j)}(\varpi) = (-1)^{j-m} (2i \sin h)^{2j} e^{ihj} \binom{2j}{j-m} \frac{[\varpi - j + m]_{2j+1}}{[\varpi + 2m]}. \quad (3.40)$$

Eq.3.38 may be re-written as

$$(J_1, M_1; J_2, M_2|J) \sim \sqrt{\frac{e^{ih(J_1+J_2-J+M_1)} e^{i\pi(J_1+M_1)}}{C_{J-J_2}^{(J_1)}(\varpi_0 + 2J_2)}} |J_1, \varpi_0 + 2J_2\rangle_{-M_1}^{J-J_2}. \quad (3.41)$$

¹¹ $\binom{n}{p}$ denotes the q-deformed binomial coefficients $\binom{n}{p} \equiv [n]!/[p]![n-p]!$

This is consistent with the orthogonality of the C.G. coefficients:

$$\sum_{M_1} \left\{ (J_1, M_1; \frac{\varpi - \varpi_0}{2}, M_2 | \frac{\varpi + 2m - \varpi_0}{2}) \times \right. \\ \left. (J_1, M_1; \frac{\varpi - \varpi_0}{2}, M_2 | \frac{\varpi + 2p - \varpi_0}{2}) \right\} = \delta_{m,p}. \quad (3.42)$$

In the limit $M_2 \rightarrow \infty$ this gives

$$\sum_{M_1} (-1)^{J_1+M_1} e^{ih(J_1+M_1)} |J_1, \varpi\rangle_{-M_1}^m |J_1, \varpi\rangle_{-M_1}^p = \delta_{m,p} C_m^{(J_1)}(\varpi) e^{ihm}. \quad (3.43)$$

This is equivalent to Eq.3.39 since $|J_1, \varpi + 2p\rangle_{-M_1}^p = e^{-ihp} |J_1, \varpi\rangle_{-M_1}^p$. Finally we use the defining relation for the 6-j symbols (Eq.2.62, or 3.13), that is,

$$\sum_{j_{23}} (j_2, \mu_2; j_3, \mu_3 | j_{23}) (j_1, \mu_1; j_{23}, \mu_{23} | j_{123}) \left\{ \begin{matrix} j_1 & j_2 \\ j_3 & j_{123} \end{matrix} \middle| \begin{matrix} j_{12} \\ j_{23} \end{matrix} \right\} = \\ (j_1, \mu_1; j_2, \mu_2 | j_{12}) (j_{12}, \mu_{12}; j_3, \mu_3 | j_{123}). \quad (3.44)$$

For the present purpose, it is convenient to let

$$j_2 = J_1, \mu_2 = -M_1, j_3 = \frac{\varpi - \varpi_0}{2}, j_{23} = \frac{\varpi - \varpi_0 + 2m_1}{2}, \\ j_1 = J_2, \mu_1 = -M_2, j_{123} = \frac{\varpi - \varpi_0 + 2m_1 + 2m_2}{2}. \quad (3.45)$$

In the limit one gets

$$\sum_{m_1+m_2=m_{12}} |J_1, \varpi\rangle_{M_1}^{m_1} |J_2, \varpi + 2m_1\rangle_{M_2}^{m_2} \sqrt{\frac{e^{ih(J_1-m_1+J_2-m_2)}}{C_{m_1}^{(J_1)}(\varpi) C_{m_2}^{(J_2)}(\varpi + 2m_1)}} \times \\ e^{i\pi(J_1+J_2)/2} \left\{ \begin{matrix} J_2 & J_1 \\ \frac{\varpi - \varpi_0}{2} & \frac{\varpi + 2m_1 + 2m_2 - \varpi_0}{2} \end{matrix} \middle| \begin{matrix} J_{12} \\ \frac{\varpi + 2m_1 - \varpi_0}{2} \end{matrix} \right\} = \\ (J_1, M_1; J_2, M_2 | J_{12}) |J_{12}, \varpi\rangle_{M_1+M_2}^{m_{12}} \sqrt{\frac{e^{ih(J_{12}-m_{12})} e^{i\pi J_{12}}}{C_{m_{12}}^{(J_{12})}(\varpi)}} \quad (3.46)$$

where we used that fact that

$$(J_2, -M_2; J_1, -M_1 | J_{12}) = (J_1, M_1; J_2, M_2 | J_{12}). \quad (3.47)$$

This relation may be verified on the explicit expression Eq.3.27. Eq.3.46 is the generalization of Eq.3.8, derived in Appendix C. Indeed, starting from the fusing algebra of the ξ fields (Eq.3.15), and taking account of the relationship between ξ and ψ fields (Eq.3.2), one concludes that the fusing of the ψ fields is of the form¹²

$$\psi_{m_1}^{(J_1)} \psi_{m_2}^{(J_2)} = \sum_{J, \{\nu\}} g_{J_1, J_2}^J N \left| \begin{matrix} J_1 & J_2 & J \\ m_1 & m_2 & m_1+m_2 \end{matrix} ; \varpi \right| \psi_{m_1+m_2}^{(J, \{\nu\})} < \varpi_J, \{\nu\} | V_{J_2-J}^{(J_1)} | \varpi_{J_2} >, \quad (3.48)$$

¹²already announced on Eq.3.16.

where

$$N \left| \begin{smallmatrix} J_1 & J_2 & J_{12} \\ m_1 & m_2 & m_1+m_2 \end{smallmatrix}; \varpi \right| = \sqrt{\frac{e^{i\pi(J_{12}-J_1-J_2)} C_{m_{12}}^{(J_{12})}(\varpi)}{C_{m_1}^{(J_1)}(\varpi) C_{m_2}^{(J_2)}(\varpi + 2m_1)}} \times e^{ih(J_1+J_2-J_{12})/2} \left\{ \begin{array}{cc|c} J_2 & J_1 & J_{12} \\ \frac{\varpi-\varpi_0}{2} & \frac{\varpi+2m_1+2m_2-\varpi_0}{2} & \frac{\varpi+2m_1-\varpi_0}{2} \end{array} \right\}. \quad (3.49)$$

Our last point is a cross check of the whole section. We re-derive the fusing matrix of the ψ fields by starting from that of the V fields (Eq.2.51), and applying the transformation Eq.3.1. The result reads

$$\begin{aligned} & \langle \varpi_{J_{123}} \{ \nu_{123} \} | \psi_{J_{23}-J_{123}}^{(J_1)} \psi_{J_3-J_{23}}^{(J_2)} | \varpi_{J_3} \{ \nu_3 \} \rangle = \\ & \sum_{J_{12}=|J_1-J_2|}^{J_1+J_2} \frac{g_{J_1 J_2}^{J_{12}} g_{J_{12} J_3}^{J_{123}} E_{J_{23}-J_{123}}^{(J_1)}(\varpi_{J_{123}}) E_{J_3-J_{23}}^{(J_2)}(\varpi_{J_{23}})}{g_{J_2 J_3}^{J_{23}} g_{J_1 J_{23}}^{J_{123}} E_{J_2-J_{123}}^{(J_{12})}(\varpi_{J_{123}})} \left\{ \begin{array}{cc|c} J_1 & J_2 & J_{12} \\ J_3 & J_{123} & J_{23} \end{array} \right\} \times \\ & \sum_{\{ \nu_{12} \}} \langle \varpi_{J_{123}} \{ \nu_{123} \} | V_{J_3-J_{123}}^{(J_{12}, \{ \nu_{12} \})} | \varpi_{J_3} \{ \nu_3 \} \rangle \langle \varpi_{J_{12}}, \{ \nu_{12} \} | V_{J_2-J_{12}}^{(J_1)} | \varpi_{J_2} \rangle. \end{aligned} \quad (3.50)$$

Comparing with Eq.3.49, we conclude that

$$\begin{aligned} \frac{g_{J_{12} J_3}^{J_{123}}}{g_{J_2 J_3}^{J_{23}} g_{J_1 J_{23}}^{J_{123}}} &= e^{-ih(J_1+J_2-J_{12})/2} \sqrt{\frac{C_{J_3-J_{123}}^{(J_{12})}(\varpi_{J_{123}}) e^{i\pi(J_{12}-J_1-J_2)}}{C_{J_{23}-J_{123}}^{(J_1)}(\varpi_{J_{123}}) C_{J_3-J_{23}}^{(J_2)}(\varpi_{J_{23}})}} \times \\ & \frac{E_{J_3-J_{23}}^{(J_{12})}(\varpi_{J_{23}})}{E_{J_{23}-J_{123}}^{(J_1)}(\varpi_{J_{123}}) E_{J_3-J_{23}}^{(J_2)}(\varpi_{J_{23}})}. \end{aligned} \quad (3.51)$$

Thus, we should have

$$g_{J_2 J_3}^{J_{23}} = e^{-ih(J_2+J_3-J_{23})/2} \sqrt{e^{i\pi(J_{23}-J_2-J_3)} C_{J_3-J_{23}}^{(J_2)}(\varpi_{J_{23}}) E_{J_3-J_{23}}^{(J_2)}(\varpi_{J_{23}})} \beta_{J_2 J_3}^{J_{23}}, \quad (3.52)$$

where β is a solution of the equation

$$\beta_{J_{12} J_3}^{J_{123}} = \beta_{J_2 J_3}^{J_{23}} \beta_{J_1 J_{23}}^{J_{123}} \quad (3.53)$$

that ensures that it disappears from Eq.3.51. In this last relation, no summation over J_{23} is understood. Since the left-hand side is independent of J_1 , this must be true for the right-hand side, and it follows that, in general, β_{LJ}^K is independent from the first lower index L . This shows that the general solution of Eq.3.53 is of the form

$$\beta_J^K = f(K)/f(J) \quad (3.54)$$

The remaining unknown function $f(I)$ is determined from the condition $g_{J_2 J_3}^{J_2+J_3} = 1$, and one finally gets

$$\begin{aligned} g_{J_2 J_3}^{J_{23}} &= e^{ih(J_{23}-J_2-J_3)/2} \sqrt{\frac{e^{i\pi(J_{23}-J_2-J_3)} C_{J_3-J_{23}}^{(J_2)}(\varpi_{J_{23}}) C_{-J_2}^{(J_2)}(\varpi_{J_2})}{C_{-J_3}^{(J_3)}(\varpi_{J_3})}} \times \\ & \frac{E_{J_3-J_{23}}^{(J_2)}(\varpi_{J_{23}}) E_{-J_2}^{(J_2)}(\varpi_{J_2})}{E_{-J_3}^{(J_3)}(\varpi_{J_3})}. \end{aligned} \quad (3.55)$$

It is straightforward, but a bit lengthy, to verify that this is equivalent to our previous expression Eq.2.39.


4 THE GENERAL (3D) STRUCTURE

In the first subsection we discuss the general structure of the bootstrap equations. We shall not give all details, but rather establish the connection with earlier works[12, 11, 25, 7] where the $U_q(sl(2))$ quantum group structure was discussed in other contexts, and show the generalization brought about by the introduction of ξ fields. Moreover, in the second part, we will introduce, by analytic continuation, a suitable limit of the general scheme spelled out in the first part, where the V and ξ operator algebras coincide.


4.1 Pictorial representations

In ref.[25], quantum-group diagrams were introduced which involve two different “worlds”: the “normal” one and the “shadow” one. Adopting this terminology from now on, we are going to verify that the OPA of the V and ξ fields is in exact correspondence with these diagrams, if the ξ and V OPE’s are associated with the normal and shadow worlds respectively. At the same time we shall discuss the associated three dimensional aspect. For the V fields it is already known, since it corresponds to the quantum-group version of the Regge-calculus approach to the discrete three-dimensional gravity[29] or to the discussion of ref.[15], for instance. This case will serve as an introduction to the novel structure that comes out when V and ξ fields are considered together.

In the pictorial representations, we omit the g coefficients. Thus, we actually make use of the operator-algebra expressed in terms of the $\tilde{V}$ fields defined by Eq.2.44, that is $\mathcal{P}_{J_{12}} \tilde{V}_{J_2-J_{12}}^{J_1} \equiv g_{J_1 J_2}^{J_{12}} \mathcal{P}_{J_{12}} V_{J_2-J_{12}}^{J_1}$, instead of the V ’s. Though of great importance for operator-product expansion, the coupling constants g define a pure gauge for the polynomial equations or knot-theory viewpoints. We could draw other figures including g coefficients, to show how they cancel out of those equations, but this would be cumbersome. The basic fusing and braiding operations on the $\tilde{V}$ operators have three equivalent representations



$$= \left\{ \begin{matrix} J_1 & J_2 \\ J_3 & J_{123} \end{matrix} \middle| \begin{matrix} J_{123} \\ J_{23} \end{matrix} \right\} \text{ (fusion of } \tilde{V} \text{ operators),} \quad (4.1)$$



$$= e^{i\pi(\Delta_{J_{123}} + \Delta_{J_3} - \Delta_{J_{23}} - \Delta_{J_{13}})} \left\{ \begin{matrix} J_1 & J_3 \\ J_2 & J_{123} \end{matrix} \middle| \begin{matrix} J_{13} \\ J_{23} \end{matrix} \right\} \text{ (braiding of } \tilde{V} \text{ operators).} \quad (4.2)$$

First consider the left diagrams, and the associated Eqs.2.51, and 2.56. Apart from the $\tilde{V}$ -matrix element on the right-hand side of the fusing relation, which has no

specific representative, each operator $\tilde{V}_m^{(J)}$ is represented by a dashed line carrying the label J . The spins on the faces display the zero-modes of the Verma modules on which the $\tilde{V}_m^{(J)}$ operators act. Thus the m 's are differences between the spin-labels of the two neighbouring faces. For the braiding diagram, the spins on the edges are unchanged at crossings, and, for given J_1, J_2 , the braiding diagram has the form of a vertex of an interaction-around-the-face (IRF) model. These diagrams are two-dimensional (2D). The appearance of spins on the faces reflect the fact that the fusion and braiding properties depend upon the Verma module on which the operators act. It is easily seen that, when they are used as building blocks, the above drawings generate diagrams which have the same structure as the quantum-group ones of ref.[25] in the shadow world¹³. The polynomial equations can be viewed as link-invariance conditions. For instance,

$$(4.3)$$


gives the pentagonal relation Eq.2.43 after cancellation of the phases (with a change of indices).

The middle diagrams of figures 4.1, and 4.2 are obtained from the left ones (first arrow) by enclosing the 2D figures with extra dashed lines carrying the spin labels which were previously on the faces. In this way, one gets three-dimensional (3D) tetrahedra, with spin labels only on the edges. The right figures are obtained from the middle ones by dualisation: the face, surrounded by the edges J_a, J_b, J_c , becomes the vertex where the edges J_a, J_b, J_c join, and conversely, a vertex becomes a face. An edge joining two vertices becomes the edge between the two dual faces. There is one triangular face for each $\tilde{V}$ field, including the $\tilde{V}$ matrix-element of the fusing relation Eq.2.51. On the dualised polyhedra, the triangular inequalities give the addition rules for spins. The main point of the middle and right diagrams is that, as a consequence of the basic MS properties of the OPA, they are really 2D projections of three-dimensional diagrams which may be rotated at essentially no cost¹⁴. For instance, the relation Eq.2.26 between fusing and braiding matrices simply corresponds to the fact that they are represented by tetrahedra which may be identified after a rigid 3D rotation. We shall illustrate the general properties of the 3D diagrams on the example of the pentagonal relation. In the same way as we closed the basic figures in Eq.4.1, 4.2, the rule to go to 3D is to close the composite figure 4.3. It gives a polyhedron which has vertices with three edges only, which we call type V3E. The two-dimensional Eq.4.3 now simply corresponds to viewing the

¹³we used dashed lines to agree with the conventions of ref.[25].


¹⁴We use 6-j symbols which do not have the full tetrahedral symmetry, so that two edges should be distinguished. This will be discussed at the end of this section.

V3E polyhedron from two different angles:



$$(4.4)$$


Dualisation gives a polyhedron, with only triangular faces, which we call F3E. The polynomial equations are recovered by decomposing a F3E polyhedron into tetrahedra (this correspondence only works with F3E polyhedra, this is the reason for dualisation). In parallel with the two different fusing-braiding decompositions of each side of Eq.4.3, there are two 3D decompositions of the F3E polyhedron. This is represented in split view on the next figure, where the internal faces are hatched for clarity.




$$(4.5)$$

In general, the rule is to take a polyhedron with triangular faces and to decompose it in tetrahedra in different ways. Substituting the associated 6-j symbols yields the polynomial identities¹⁵.

In quantum-group diagrams[25], a second world was introduced – the normal one – which is represented by solid lines. In this world, the quantum numbers are spins J and magnetic numbers M both on the lines. The label M changes at the crossings. We now show how our ξ operator algebra is a realisation of this normal world. Indeed, the fusing and braiding matrices of the fields $\xi_M^{(J)}$ do not depend upon the Verma module on which they act, so that it is consistent that the quantum numbers J and M be attached to the corresponding line. Comparing Eqs.3.14, and 3.21 with the formulae given in ref.[25], one sees that one has the following 2D representation¹⁶



$$(J_1, M_1; J_2, M_2 | J_{12}) = \text{fusion of } \xi \text{ operators}, \quad (4.6)$$



$$(J_1, J_2)_{M_1' M_2'}^{M_2' M_1'} = \text{braiding of } \xi \text{ operators}, \quad (4.7)$$

¹⁵we restrict ourselves to polyhedra which are orientable surfaces.

¹⁶For Eq.3.14, the factor $g_{J_1 J_2}^{J_{12}}$ is absorbed by going from V to $\tilde{V}$ matrix element on the right-hand side.

which coincides with the corresponding quantum-group vertices of ref.[25]. Clearly, the braiding diagram Eq.4.7 should be regarded as an interaction of the vertex type. In the same way as for the $\tilde{V}$ fields, the fusing vertex has only three legs so that the $\tilde{V}$ matrix element is not represented per se. On the other hand, we have shown, using repeated fusions (Eqs.3.16-3.26) that the fusing equation also provides the transitions between $\tilde{V}$ and ξ operator-product algebras. From this viewpoint, the result of the discussion just recalled may be pictorially represented by introducing transitions between the two worlds based on the graph

$$\begin{array}{c} M_1 \\ \swarrow \\ J_1 \end{array} \begin{array}{c} M_2 \\ \searrow \\ J_2 \end{array} \begin{array}{c} M_{12} \\ \swarrow \\ J_{12} \end{array} = (J_1, M_1; J_2, M_2 | J_{12}) \text{ (from normal to shadow world),} \quad (4.8)$$

which coincides with the one introduced in ref.[25].

The polynomial equations involving $\tilde{V}$, and/or ξ fields are summarized by the link-invariance of the diagrams constructed out of the building blocks just given. For instance, the example previously given for the $\tilde{V}$ operators becomes, in the normal world,

$$\begin{array}{c} M_1 \\ \swarrow \\ J_1 \end{array} \begin{array}{c} M_2 \\ \searrow \\ J_2 \end{array} \begin{array}{c} M_3 \\ \swarrow \\ J_3 \end{array} \begin{array}{c} M_{12} \\ \swarrow \\ J_{12} \end{array} \begin{array}{c} M_{23} \\ \swarrow \\ J_{23} \end{array} \begin{array}{c} M_{1'1} \\ \swarrow \\ J_{1'1} \end{array} \begin{array}{c} M_{2'2} \\ \swarrow \\ J_{2'2} \end{array} \begin{array}{c} M_{3'3} \\ \swarrow \\ J_{3'3} \end{array} = \begin{array}{c} M_1 \\ \swarrow \\ J_1 \end{array} \begin{array}{c} M_2 \\ \searrow \\ J_2 \end{array} \begin{array}{c} M_3 \\ \swarrow \\ J_3 \end{array} \begin{array}{c} M_{2+3} \\ \swarrow \\ J_{2+3} \end{array} \begin{array}{c} M_{1'1} \\ \swarrow \\ J_{1'1} \end{array} \quad (4.9)$$

Let us establish a three-dimensional representation involving ξ operators. We propose the following

$$\begin{array}{c} M_1 \\ \swarrow \\ J_1 \end{array} \begin{array}{c} M_2 \\ \searrow \\ J_2 \end{array} \begin{array}{c} M_{12} \\ \swarrow \\ J_{12} \end{array} \rightarrow \begin{array}{c} M_1 \\ \swarrow \\ J_1 \end{array} \begin{array}{c} M_2 \\ \searrow \\ J_2 \end{array} \begin{array}{c} M_{12} \\ \swarrow \\ J_{12} \end{array} \rightarrow \begin{array}{c} M_1 \\ \swarrow \\ J_1 \end{array} \begin{array}{c} M_2 \\ \searrow \\ J_2 \end{array} \begin{array}{c} M_{12} \\ \swarrow \\ J_{12} \end{array} \quad (4.10)$$

$$\begin{array}{c} M_1 \\ \swarrow \\ J_1 \end{array} \begin{array}{c} M_2 \\ \searrow \\ J_2 \end{array} \begin{array}{c} M_{12} \\ \swarrow \\ J_{12} \end{array} \rightarrow \begin{array}{c} M_1 \\ \swarrow \\ J_1 \end{array} \begin{array}{c} M_2 \\ \searrow \\ J_2 \end{array} \begin{array}{c} M_{12} \\ \swarrow \\ J_{12} \end{array} \rightarrow \begin{array}{c} M_1 \\ \swarrow \\ J_1 \end{array} \begin{array}{c} M_2 \\ \searrow \\ J_2 \end{array} \begin{array}{c} M_{12} \\ \swarrow \\ J_{12} \end{array} \quad (4.11)$$

When the left diagrams are enclosed, we put the M 's at the edges. The surrounding lines are drawn as solid, while the lines which already existed become dashed. This ensures consistency with the tetrahedral representation of the $\tilde{V}$ operator-product

algebra given above, since dashed lines have a J label in agreement with the previous convention – contrary to the solid ones. In the dualisation, the dashed lines are transformed as before, while the M 's naturally go on the faces. These come out of two types. In the fusion, there is one face which has no M and is surrounded by three dashed lines. It represents the $\tilde{V}$ field which appears in Eq.3.14. All other faces in the two above diagrams are similar: surrounded by two solid lines with no label, and a dashed line with a J label, they carry the corresponding magnetic number M . Each of them represents a ξ field. In the dualised diagrams, the face associated with $\tilde{V}$ fields is drawn as transparent, while the ξ faces are dashed. For them we also give a top-view drawn on the lower-left quarter plane, and a left-view drawn on the upper-right quarter plane. The tetrahedral representation of the diagram Eq.4.8 is chosen so that its 3D representations agrees with the one of the fusion up to a rotation since they are given by the same 3-j symbols. Thus we let

(4.12)

The 3D aspect now summarizes the general polynomial equations. The relation displayed by the figure 4.9 is transformed into


(4.13)

The dualised polyhedron is

(4.14)

where all the faces are dashed except the face J_2, J_3, J_{23} , of the $\tilde{V}$ type, which is transparent, according with the general convention. We next draw its decomposition, altering the general convention for visibility: the external quantum number are

not written, and some external faces are made transparent



Consider in general a higher 2D diagram with one separation between a shadow and a real part. The enclosure proceeds as follows. In each world the rule is as indicated above. Concerning the separation line, one follows the prescription suggested by figure 4.12, namely, the separation line becomes solid, and thus carries no label. As a result, the higher 3D diagram before dualisation is again of the V3E type, with any number of dashed lines, and one closed loop of solid lines. Thus there are only two types of vertices: with three dashed lines, or with one dashed and two solid lines. After dualisation, one may obtain any polyhedron of the F3E type with the two kinds of faces introduced above: faces of the $\tilde{V}$ type (three dashed lines, each with a J , around a – transparent – face), and faces of the ξ type (two solid lines with no number, a dashed line with a J , and an M on the face – which is dashed). A J has to be interpreted as the length of the corresponding edge, and an M as the difference of length of the two surrounding solid edges. This gives all the spin addition rules and relations between M 's as triangular inequalities. Of course, the J value of a dashed line is common to the two adjacent faces. The polynomial equations are derived by splitting a general F3E polyhedron in tetrahedra. Since there are two types of faces, one can only obtain three types of tetrahedra. There is a first type of tetrahedron, with three $\tilde{V}$ and one ξ faces, (see figure 4.10 or 4.12), its value is the corresponding Clebsch-Gordan coefficient. The tetrahedra of the second type have four ξ faces, (see figure 4.11), the values are the corresponding R-matrix elements. The third type tetrahedron has four $\tilde{V}$ faces (see figure 4.1 or 4.2), its value is the corresponding 6-j coefficient. In the case of the third type tetrahedron, changing the orientation yields an extra phase as we already mentioned.

For completeness, we have to add that the 6-j, R-matrix or Clebsch-Gordan have less symmetries than the tetrahedra by which they are represented, and therefore that these symmetries must be broken by adding extra characteristics to the tetrahedra, so that the correspondence be one-to-one. We give them briefly. First, the faces of the tetrahedra must be either 'incoming' or 'outgoing'. Each tetrahedron has two incoming faces (the faces J_1, J_{23}, J_{123} and J_2, J_3, J_{23} in figures 4.1 and 4.2, the faces M_1 and M_2 in figures 4.10, 4.11 and 4.12) and two outgoing ones (the two other ones). When splitting a composite polyhedron in tetrahedra, the internal faces, common to two tetrahedra, are outgoing for one tetrahedron and incoming for the other one. This rule allows to single out the two particular J of the non-symmetric non-RW 6-j (the J between the two incoming faces, and the J between

the two outgoing ones), to distinguish M_1 and M_2 from M_{12} for the C-G, and to distinguish between the M_i 's and M'_i 's of the R-matrix. It is clear that the incoming faces represent the operators on which the fusing or braiding are performed, and the outgoing faces the resulting operators. Secondly, the M_i 's on the faces are oriented quantities. Like the m_i 's, they should be considered as differences of lengths of the two solid lines surrounding the face M_i , which thus must be supplemented by an ordering. This ordering must always be from left to right (for instance) on the 2D projection. If we rotate the tetrahedron of figure 4.11 representing $(J_1, J_2)_{M_1 M_2}^{M'_1 M'_2}$, by π around a vertical axis in the plane of the page, this exchanges 1 and 2, but the orderings as well. The signs of the M_i 's must therefore be changed to restore the left-right ordering of the 2D projection. The resulting value is then $(J_2, J_1)_{-M_2 -M_1}^{-M'_1 -M'_2}$, which is indeed equal to $(J_1, J_2)_{M_1 M_2}^{M'_1 M'_2}$. So, these tetrahedra with two outgoing and two incoming faces, and oriented M_i 's, are in one-to-one correspondence with the 6-j, R-matrix or Clebsch-Gordan coefficients.

4.2 ξ as limit of $\tilde{V}$

In part 4.1, we showed that the V braiding diagrams could be interpreted as IRF model vertices and the ξ braiding diagrams as vertex model interactions. Witten showed that one could get vertex models as limit of IRF models[15], when letting the spins on the faces go to infinity with fixed differences. In this part we shall apply this method directly to our operators. The spins on the faces are the ones corresponding to zero-modes. We shall denote by I_ϖ the left-most one (it is defined by $\varpi \equiv \varpi_0 + 2I_\varpi$), and let ϖ go to infinity. Hence, getting a vertex model as limit of an IRF model is equivalent to obtaining ξ operators as limits of $\tilde{V}$ operators. More precisely, let us prove that

$$\lim_{\varpi \rightarrow \infty} \mathcal{P}_{I_\varpi} \tilde{V}_m^{(J)} / \beta_{I_\varpi+m}^{I_\varpi} = (2i)^m e^{-ihm} \xi_m^{(J)} \quad (4.16)$$

The role of the β coefficients is to remove the Γ functions which would have no well defined limit. They will cancel out of the braiding or fusing identities thanks to their property Eq.3.53. $\mathcal{P}_{I_\varpi}$ is the projector on the Verma module of spin I_ϖ defined in Eq.2.23. Moreover, we have to give an imaginary part to I_ϖ (or ϖ) so that the limit $e^{\pm i h \varpi}$ be well defined: we choose a negative imaginary part, which makes $e^{-i h \varpi}$ go to zero and $e^{i h \varpi}$ to infinity.

Using the expression Eq.3.52 of g and $\tilde{V}_m^{(J)} = (g_{J I_\varpi+m}^{I_\varpi} / E_m^{(J)}(\varpi)) \sum_M (J, \varpi |_m^M \xi_M^{(J)})$ where $(J, \varpi |_m^M$ is the inverse matrix of $|J, \varpi \rangle_M^m$, Eq.4.16 is equivalent to

$$\lim_{\varpi \rightarrow \infty} \mathcal{P}_{I_\varpi} e^{ih(J+m)/2} \sqrt{e^{i\pi(J+m)} C_m^{(J)}(\varpi)} \sum_M (J, \varpi |_m^M \xi_M^{(J)}) = (2i)^m e^{-ihm} \xi_m^{(J)}. \quad (4.17)$$

Let us prove now that the matrix $|J, \varpi \rangle_M^m$ has a leading term proportional to the identity matrix when $\varpi \rightarrow \infty$, and consequently, such is the leading term of its inverse matrix. Recall Eq.3.35

$$|J, \varpi \rangle_M^m = \sqrt{\binom{2J}{J+M}} e^{ihm/2} \times \sum_{(J-M+m-t)/2 \text{ integer}} e^{iht(\varpi+m)} \binom{J-M}{(J-M+m-t)/2} \binom{J+M}{(J+M+m+t)/2}. \quad (4.18)$$

It is a polynomial in $e^{ih\varpi}$. As $e^{ih\varpi}$ has an infinite limit, the maximum value of t dominates in the sum. The boundaries on t are given by the condition that the q -factorials arguments be positive. Thus, we get

$$\lim_{\varpi \rightarrow \infty} e^{-ihJ\varpi} |J, \varpi\rangle_M^m = \sqrt{\binom{2J}{J+M}} e^{ihM(J+1/2)} \delta_{m,M}. \quad (4.19)$$

This gives the limit of the inverse matrix directly. In view of Eq.4.17 we also need the limit of $C_m^{(J)}(\varpi)$. Since the complex exponential with positive argument is dominant in $\sin(h\varpi)$, we get

$$\lim_{\varpi \rightarrow \infty} e^{-ih2J\varpi} C_m^{(J)}(\varpi) = (-1)^{-J-m} e^{ihJ} \binom{2J}{J-m} e^{ih(2J-1)m}, \quad (4.20)$$

which leads to Eq.4.16.

Now, we take this limit $\varpi \rightarrow \infty$ in the braiding or fusing equations. Begin with the fusing Eq.2.51. Let us write it down in term of three $\tilde{V}$ and one V

$$\begin{aligned} \mathcal{P}_{I_\varpi} \tilde{V}_{m_1}^{(J_1)} \tilde{V}_{m_2}^{(J_2)} &= \sum_{J_{12}=|J_1-J_2|}^{J_1+J_2} g_{J_1 J_2}^{J_{12}} \left\{ \begin{matrix} J_1 & J_2 \\ I_\varpi+m_1+m_2 & I_\varpi \end{matrix} \middle| \begin{matrix} J_{12} \\ I_\varpi+m_1 \end{matrix} \right\} \times \\ &\sum_{\{\nu_{12}\}} \mathcal{P}_{I_\varpi} \tilde{V}_{m_1+m_2}^{(J_{12}, \{\nu_{12}\})} < \varpi_{J_{12}}, \{\nu_{12}\} | V_{J_2-J_{12}}^{(J_1)} | \varpi_{J_2} >. \end{aligned} \quad (4.21)$$

The β coefficients introduced by Eq.4.16 cancel out thanks to the property Eq.3.53, and, in the limit, we do get the fusing Eq.3.14 of the ξ operators, provided that

$$\lim_{\varpi \rightarrow \infty} \left\{ \begin{matrix} J_1 & J_2 \\ I_\varpi+m_1+m_2 & I_\varpi \end{matrix} \middle| \begin{matrix} J_{12} \\ I_\varpi+m_1 \end{matrix} \right\} = (J_1, m_1; J_2, m_2 | J_{12}). \quad (4.22)$$

Here again, we have to give an imaginary part to ϖ so that the limits of the complex exponentials be well defined. But the 6-j coefficients are only defined for positive half-integer spins. So, we have to extend this definition to non integer I_ϖ before going to the limit. Such was not the case for the limit of $\mathcal{P}_{I_\varpi} \tilde{V}_m^{(J)}$ considered above, since everything was defined for any ϖ . This gives us the condition that our extended definition of the 6-j must be coherent with the limit of the $\tilde{V}$, i.e. it must obey Eq.4.22.

The expression of the 6-j coefficients is given in Eq.2.54. The ambiguity of the extension lies in the range of summation. For half-integer spins the boundaries are given by the condition that the arguments of the q -factorial in the denominator must not be negative integers. For half integer I_ϖ the initial definition is strictly equivalent to the following one, obtained by the change of index $\mu = z - 2I_\varpi - J_{12} - m_1 - m_2$

$$\begin{aligned} \left\{ \begin{matrix} J_1 & J_2 \\ I_\varpi+m_1+m_2 & I_\varpi \end{matrix} \middle| \begin{matrix} J_{12} \\ I_\varpi+m_1 \end{matrix} \right\} &= (-1)^{J_{12}-J_1-J_2} \sqrt{[2J_{12}+1][2I_\varpi+2m_1+1]} \Delta(J_1, J_2, J_{12}) \times \\ &\Delta(J_1, I_\varpi, I_\varpi+m_1) \Delta(I_\varpi+m_1+m_2, I_\varpi, J_{12}) \Delta(I_\varpi+m_1+m_2, J_2, I_\varpi+m_1) \sum_{\mu \text{ integer}} (-1)^\mu \times \\ &[2I_\varpi+J_{12}+m_1+m_2+\mu+1]! \left[[2I_\varpi-J_1-J_2+m_1+m_2+\mu]! [J_{12}-J_1+m_2+\mu]! \times \right. \\ &\left. [J_{12}-J_2-m_1+\mu]! [\mu]! [J_1+J_2-J_{12}-\mu]! [J_1+m_1-\mu]! [J_2-m_2-\mu]! \right]^{-1}. \end{aligned} \quad (4.23)$$

But, when we give an imaginary part to I_ϖ , the q-factorials with a I_ϖ have complex arguments and yield no restriction. Such is the case of the last six factorials of the sum with the definition of Eq.2.54, and, of the first two with the definition of Eq.4.23. The first definition rapidly appears inadequate as it yields no upper boundary for z . The second definition (Eq.4.23) leads to well defined boundaries for the index μ and to a finite limit thanks to

$$\frac{[\alpha + 2I_\varpi]!}{[\beta + 2I_\varpi]!} \sim (2i \sin h)^{\beta-\alpha} e^{ih(\alpha-\beta)2I_\varpi} e^{ih(\alpha-\beta)(\alpha+\beta+1)/2} \quad (4.24)$$

for $\varpi \rightarrow \infty$ with negative imaginary part, similar to Eq.3.33 which was for positive imaginary part.

We use this limit in the sum and in the Δ prefactors of Eq.4.23, and recognize the expression of the Clebsch-Gordan coefficient given in Eq.3.27. This justifies our extended definition of the 6-j.

Pictorially¹⁷, this reads

$$(4.25)$$

We come now to the case of braiding. It works like fusing. From Eq.4.16 and the properties of the β coefficients Eq.3.53, we see that we only have to prove that when J goes to infinity with negative imaginary part

$$\lim_{\varpi \rightarrow \infty} \left\{ \begin{matrix} J_1 & I_\varpi + m \\ J_2 & I_\varpi \end{matrix} \middle| \begin{matrix} I_\varpi + m'_2 \\ I_\varpi + m_1 \end{matrix} \right\} e^{i\pi(\Delta_{I_\varpi+m} + \Delta_{I_\varpi} - \Delta_{I_\varpi+m'_2} - \Delta_{I_\varpi+m_1})} = (J_1, J_2)_{m_1 m_2}^{m'_1 m'_2} \quad (4.26)$$

and

$$\lim_{\varpi \rightarrow \infty} \left\{ \begin{matrix} J_1 & I_\varpi + m \\ J_2 & I_\varpi \end{matrix} \middle| \begin{matrix} I_\varpi + m'_2 \\ I_\varpi + m_1 \end{matrix} \right\} e^{-i\pi(\Delta_{I_\varpi+m} + \Delta_{I_\varpi} - \Delta_{I_\varpi+m'_2} - \Delta_{I_\varpi+m_1})} = \overline{(J_1, J_2)_{m_1 m_2}^{m'_1 m'_2}} \quad (4.27)$$

where $m \equiv m_1 + m_2 = m'_1 + m'_2$ and (J_1, J_2) (resp. $\overline{(J_1, J_2)}$) is the universal R-matrix R (resp. $\overline{R}$), following the conventions of ref[19] (see Eq.3.22). We only deal in details with the case of the R-matrix R . We compute its matrix element from its universal form given in Eq.3.23

$$(J_1, J_2)_{m_1 m_2}^{m'_1 m'_2} = e^{-2ihm_1 m_2} \frac{(-1)^n (2i \sin(h))^n e^{ihn(n+1)}}{[n]!} e^{-ihn m_1} e^{ihn m_2} \times \sqrt{\frac{[J_1 + m_1]! [J_1 - m'_1]! [J_2 - m_2]! [J_2 + m'_2]!}{[J_1 - m_1]! [J_1 + m'_1]! [J_2 + m_2]! [J_2 - m'_2]!}}, \quad (4.28)$$

for $n \equiv m'_2 - m_2 \geq 0$, and 0 otherwise. It is an upper triangular matrix.

To begin with, we examine the behavior of the 6-j coefficient when ϖ goes to infinity with a negative imaginary part. This time, the suitable change of summation index to define the 6-j for non-integer spins is $x = 2I_\varpi + J_1 + J_2 + m_1 + m_2 - z$

¹⁷for a better readability, we omit the index ϖ of I_ϖ in the drawings

(basically, this is the same definition of the extension as before, which amounts to taking $z - 2I_\varpi$ integer, the extra integer shifts by $J_i + m_i$ being trivial). Here, only the first and eighth factorials of the sum have infinite arguments. After using Eq.4.24, we get

$$\begin{aligned} \left\{ \begin{matrix} J_1 & I_\varpi+m \\ J_2 & I_\varpi \end{matrix} \middle| \begin{matrix} I_\varpi+m'_2 \\ I_\varpi+m_1 \end{matrix} \right\} &\sim e^{-ih2I_\varpi n} e^{ih(m_1+m'_2+1)} \sqrt{[J_1+m_1]![J_1-m_1]![J_1+m'_1]![J_1-m'_1]} \\ &\times \sqrt{[J_2+m_2]![J_2-m_2]![J_2+m'_2]![J_2-m'_2]} (2i \sin(h))^n e^{-ih/2(1+n)(m+m_1+m'_2+2)} \times \\ &\sum_{x \text{ integer}} \frac{e^{-4ihI_\varpi x} (2i \sin(h))^{2x} e^{-ihx(m+m_1+M'_2+2)} (-1)^x}{[J_1-m_1-x]![J_1+m'_1-x]![J_2+m_2-x]![J_2-m'_2-x]! [x]! [n+x]!}. \end{aligned} \quad (4.29)$$

As $e^{-ihI_\varpi} \rightarrow 0$, due to the factor $e^{-4ihI_\varpi x}$ the term of the sum with x minimal will be dominant. The lower boundary for x is given by $[x]!$ and $[n+x]!$: for $n \geq 0$, the minimal allowed x is 0, and for $n \leq 0$, it is n . Hence, summarizing both cases, the dominant behavior of the 6-j is $e^{-ih2I_\varpi |n|}$.

Then, the extra factor due to $e^{\pm i\pi(\Delta_{I_\varpi+m}+\Delta_{I_\varpi}-\Delta_{I_\varpi+m'_2}-\Delta_{I_\varpi+m_1})}$, which is $e^{\pm ih2I_\varpi n} \times$ (finite term), must be taken into account. Altogether, it gives $e^{ih2I_\varpi(\pm n-|n|)}$, i.e. $e^{-ih4I_\varpi |n|}$ when $n \leq 0$, and 1 when $n \geq 0$, in the upper case(+). So, when $e^{-ih2I_\varpi} \rightarrow 0$, the limit is an upper triangular matrix, as it should (In the lower case(-) it gives 1 when $n \leq 0$, and $e^{-ih4I_\varpi |n|}$ when $n \geq 0$ and so, a lower triangular matrix in the limit).

It is then straightforward to check that the limit agrees with Eq.4.28, and this terminates the proof of Eq.4.26.

The pictorial representation of this is

$$(4.30)$$

We can use these limits on more complex braiding-fusing identities of $\tilde{V}$ fields, thereby proving the same identities for ξ fields. For instance, we have the following limit

$$(4.31)$$

where we did not write the spins J_i on the lines as they are not affected by the limit, and where the missing m_i on the r.h.s can be deduced from conservation of $\sum_i m_i$ at each vertex. Making this limit on both sides of identities in the shadow world yields identities in the normal world.

We can as well make a limit only on a part of the figure. Eq.4.22 gives as well

$$\frac{e^{-i\pi((\Delta_{I+m_1+m_2}-\Delta_{J_{12}}))}}{e^{-i\pi(\Delta_{I+m_1}-\Delta_{J_1}))}} \times \begin{array}{c} I+m_1 \quad J_2 \\ \diagup \quad \diagdown \\ J_1 \quad I+m_1+m_2 \\ \diagdown \quad \diagup \\ J_{12} \end{array} \xrightarrow{I \rightarrow \infty} \begin{array}{c} m_1 \quad m_2 \\ \diagup \quad \diagdown \\ J_1 \quad J_{12} \\ \diagdown \quad \diagup \\ m_1+m_2 \end{array} \quad (4.32)$$

which gives on a composite figure

$$\frac{e^{-i\pi(\Delta_{I+m_1+m_2+m_3+m_4}-\Delta_{J_c})}}{e^{-i\pi(\Delta_{I+m_1}-\Delta_{J_a})}} \times \begin{array}{c} I+m_1 \quad I+m_1+m_2 \\ \diagup \quad \diagdown \\ I+m_1+m_2 \quad I+m_1+m_2+m_3 \\ \diagdown \quad \diagup \\ J_a \quad J_b \quad I+m_1+m_2+m_3+m_4 \\ \diagdown \quad \diagup \\ J_c \end{array} \xrightarrow{I \rightarrow \infty} \begin{array}{c} m_1 \quad m_2 \quad m_3 \quad m_4 \\ \diagup \quad \diagdown \\ J_a \quad J_b \quad J_c \end{array} \quad (4.33)$$

The extra phases in Eq.4.32 cancel out between two neighbouring basic figures all along the wavy line, and only the exterior ones remain as in Eq.4.33. Hence, they do not spoil equalities between transformed figures since the exterior spins of both sides are the same.

To conclude with this part, we show how this limit can be understood in the case of the 3D representation by V3E polyhedra. As an example, we close the 2D figures 4.31 and 4.33 and get the 3D representation

$$\begin{array}{c} I+m_1 \quad I+m_1+m_2 \\ \diagup \quad \diagdown \\ I+m_1+m_2 \quad I+m_1+m_2+m_3 \\ \diagdown \quad \diagup \\ I+m_1+m_2+m_3+m_4 \\ \diagdown \quad \diagup \\ I+m_6 \quad I+m_6+m_7 \\ \diagdown \quad \diagup \\ I \end{array} \xrightarrow{I \rightarrow \infty} \begin{array}{c} m_1 \quad m_2 \quad m_3 \quad m_4 \\ \diagup \quad \diagdown \\ m_6 \quad m_7 \quad m_6+m_7-m_1-m_2-m_3-m_4 \end{array} \quad (4.34)$$

and

$$\frac{e^{-i\pi(\Delta_{I+m_1+m_2+m_3+m_4}-\Delta_{J_c})}}{e^{-i\pi(\Delta_{I+m_1}-\Delta_{J_a})}} \times \begin{array}{c} I+m_1 \quad I+m_1+m_2 \\ \diagup \quad \diagdown \\ I+m_1+m_2 \quad I+m_1+m_2+m_3 \\ \diagdown \quad \diagup \\ J_a \quad J_b \quad I+m_1+m_2+m_3+m_4 \\ \diagdown \quad \diagup \\ J_c \end{array} \xrightarrow{I \rightarrow \infty} \begin{array}{c} m_1 \quad m_2 \quad m_3 \quad m_4 \\ \diagup \quad \diagdown \\ J_a \quad J_b \quad J_c \end{array} \quad (4.35)$$

where again the spins on the internal lines, which are unchanged by the limit have not been written down.

We see on these examples that the partial or global limits are not fundamentally different. On the V3E polyhedra, the normal world is obtained from the shadow world by adding I to all the edges of a arbitrarily chosen closed loop, and then letting I go to infinity. In this limit, only the differences between the values on the edges remain finite and relevant; they go on the vertices. When projecting in two

dimensions, the part of the drawing inside the projection of the closed loop naturally becomes the normal world, as the part outside becomes the shadow world. Hence, depending on whether the closed loop is at the exterior of the drawing or not, there is only a normal world (3D in figure 4.34, 2D in figure 4.31) or both worlds (3D in figure 4.35, 2D in figure 4.33), but fundamentally, this is not different.

Following the remark of the end of part 4.1, we note that we need to orient the M_i 's at the vertices of the V3E polyhedra (and then on the faces of the F3E polyhedra) in order to know how to order the difference of the two neighbouring spins in the infinite limit.

5 FUSION RULES FOR $\xi_{M\widehat{M}}^{(J\widehat{J})}$ OPERATORS

In the previous sections, we have been able to determine the coupling constants of the fusion rules for the operators connected with the quantum group parameter h , by assuming that the general scheme of Moore and Seiberg holds. In this section, we deal with the most general operators $V_{M\widehat{M}}^{(J\widehat{J})}$, and $\xi_{M\widehat{M}}^{(J\widehat{J})}$ that involve the two quantum group parameters h and $\widehat{h}$. It is straightforward to extend the full discussion just given to this case, since the fusing and braiding matrices of the $V_M^{(J)}$ (or equivalently of the $\xi_M^{(J)}$) with the $V_{\widehat{M}}^{(\widehat{J})}$ (or equivalently with the $\xi_{\widehat{M}}^{(\widehat{J})}$) operators are simple phases. We shall not do it explicitly to keep the length of this paper within reasonable limits. We shall only determine the coupling constants by using a short cut that only assumes that the short-distance expansion of the product of two fields ξ do not depend on ϖ , and computes its matrix element between the state $|\varpi_0\rangle$ and the appropriate highest-weight-state bra, in two equivalent ways. In order to illustrate the method, let us first rederive the expression Eq.2.39 for $g_{J_1 J_2}^{J_{12}}$. We shall later on extend the method to the fusion of the $\xi_{M\widehat{M}}^{(J\widehat{J})}$ fields.

Let us compute $\langle \varpi_{J_3} | \xi_{M_1}^{(J_1)}(z) \xi_{M_2}^{(J_2)}(0) | \varpi_0 \rangle$ in two different ways

(1) directly using the relation between $\xi_M^{(J)}$ and $V_M^{(J)}$ as well as equation (2.10)

$$\begin{aligned} & \langle \varpi_{J_3} | \xi_{M_1}^{(J_1)}(z) \xi_{M_2}^{(J_2)}(0) | \varpi_0 \rangle = \\ & E_{J_2-J_3}^{(J_1)}(\varpi_{J_3}) E_{-J_2}^{(J_2)}(\varpi_{J_2}) | J_1, \varpi_{J_3} \rangle_{M_1}^{J_2-J_3} | J_2, \varpi_{J_2} \rangle_{M_2}^{-J_2} z^{(\Delta_3-\Delta_1-\Delta_2)}; \end{aligned} \quad (5.1)$$

(2) using first the short distance expansion

$$\xi_{M_1}^{(J_1)}(z) \xi_{M_2}^{(J_2)}(0) \simeq \sum_{J_{12}} z^{\Delta_{12}-\Delta_1-\Delta_2} [G_{M_1 M_2 M_1+M_2}^{J_1 J_2 J_{12}} \xi_{M_1+M_2}^{J_{12}}(0) + \dots] \quad (5.2)$$

where G is to be determined. Then

$$\begin{aligned} & \langle \varpi_{J_3} | \xi_{M_1}^{(J_1)}(z) \xi_{M_2}^{(J_2)}(0) | \varpi_0 \rangle = \\ & G_{M_1 M_2 M_1+M_2}^{J_1 J_2 J_3} E_{-J_3}^{(J_3)}(\varpi_{J_3}) | J_3, \varpi_{J_3} \rangle_{M_1+M_2}^{-J_3} z^{(\Delta_3-\Delta_1-\Delta_2)} + \dots \end{aligned} \quad (5.3)$$

One gets immediately

$$G_{M_1 M_2 M_1+M_2}^{J_1 J_2 J_3} = \frac{E_{J_2-J_3}^{(J_1)}(\varpi_{J_3}) E_{-J_2}^{(J_2)}(\varpi_{J_2}) | J_1, \varpi_{J_3} \rangle_{M_1}^{J_2-J_3} | J_2, \varpi_{J_2} \rangle_{M_2}^{-J_2}}{E_{-J_3}^{(J_3)}(\varpi_{J_3}) | J_3, \varpi_{J_3} \rangle_{M_1+M_2}^{-J_3}} \quad (5.4)$$

Applying Eq.3.46 with $m_{12} = -J_3$, $\varpi = \varpi_{J_3}$ and noting that due to triangular inequalities the non symmetric 6-j coefficient $\left\{ \begin{matrix} J_2 & J_1 \\ J_3 & 0 \end{matrix} \middle| \begin{matrix} J_3 \\ J_3+m_1 \end{matrix} \right\}$ is different from 0 only for $m_1 = J_2 - J_3$ we get

$$\frac{|J_1, \varpi_{J_3}\rangle_{M_1}^{J_2-J_3} |J_2, \varpi_{J_2}\rangle_{M_2}^{-J_2}}{|J_3, \varpi_{J_3}\rangle_{M_1+M_2}^{-J_3}} = e^{[ih(J_3-J_1-J_2)+i\pi(J_3-J_1-J_2)]/2} \times \sqrt{\frac{C_{J_2-J_3}^{(J_1)}(\varpi_{J_3}) C_{-J_2}^{(J_2)}(\varpi_{J_2})}{C_{-J_3}^{(J_3)}(\varpi_{J_3})}} (J_1 M_1 J_2 M_2 | J_3) \quad (5.5)$$

As expected, this gives

$$G_{M_1 M_2 M_1+M_2}^{J_1 J_2 J_{12}} = g_{J_1 J_2}^{J_{12}} (J_1 M_1 J_2 M_2 | J_{12}) \quad (5.6)$$

where $g_{J_1 J_2}^{J_{12}}$ is given by Eq.3.55. It is worth to note that, instead of using the short distance expansion Eq.5.2, it is consistent to use the exact Eq.3.14 and we get the same result due to the fact that $\langle \varpi_{J_3} | V_{M_1+M_2}^{(J_{12}, \nu)}(0) | \varpi_0 \rangle$ is different from 0 only for the highest weight operator $V_{M_1+M_2}^{(J_{12})} (\nu = 0)$.

The same method applies exactly to the fusion of the $\xi_{M\hat{M}}^{(J\hat{J})}$ fields using now[19]

$$\xi_{M\hat{M}}^{(J\hat{J})} = e^{4i\pi J\hat{J}} e^{-i\pi(M\hat{J}+\hat{M}J)} \sum_{m, \hat{m}} |J, \varpi\rangle_M^m | \hat{J}, \hat{\varpi} \rangle_{\hat{M}}^{\hat{m}} E_{m\hat{m}}^{(J\hat{J})}(\varpi) V_{m\hat{m}}^{(J\hat{J})} \quad (5.7)$$

and

$$|J, \varpi + 2\hat{\ell} \frac{\pi}{h}\rangle_M^m = e^{2i\pi \hat{\ell} m} e^{-2i\pi \hat{\ell} (J-M)} |J, \varpi\rangle_M^m. \quad (5.8)$$

The short distance expansion of the product of two $\xi_{M\hat{M}}^{J\hat{J}}$ fields is written as

$$\xi_{M_1 \hat{M}_1}^{(J_1 \hat{J}_1)}(z) \xi_{M_2 \hat{M}_2}^{(J_2 \hat{J}_2)}(0) \simeq \sum_{J_3, \hat{J}_3} z^{\Delta_{J_3 \hat{J}_3} - \Delta_{J_1 \hat{J}_1} - \Delta_{J_2 \hat{J}_2}} \times [G_{M_1 \hat{M}_1 M_2 \hat{M}_2 M_1+M_2 \hat{M}_1+\hat{M}_2}^{J_1 \hat{J}_1 J_2 \hat{J}_2 J_3 \hat{J}_3} \xi_{M_1+M_2 \hat{M}_1+\hat{M}_2}^{(J_3 \hat{J}_3)}(0) + \dots]. \quad (5.9)$$

The final result reads

$$G_{M_1 \hat{M}_1 M_2 \hat{M}_2 M_1+M_2 \hat{M}_1+\hat{M}_2}^{J_1 \hat{J}_1 J_2 \hat{J}_2 J_3 \hat{J}_3} = g_{J_1 \hat{J}_1 J_2 \hat{J}_2}^{J_3 \hat{J}_3} (J_1 M_1 J_2 M_2 | J_3) (\hat{J}_1 \hat{M}_1 \hat{J}_2 \hat{M}_2 | \hat{J}_3) \times e^{i\pi[M_1 \hat{J}_2 - M_2 \hat{J}_1 + \hat{M}_1 J_2 - \hat{M}_2 J_1 + (M_1 - M_2)(\hat{J}_3 - \hat{J}_1 - \hat{J}_2) + (\hat{M}_1 - \hat{M}_2)(J_3 - J_1 - J_2)]} \quad (5.10)$$

with

$$g_{J_1 \hat{J}_1 J_2 \hat{J}_2}^{J_3 \hat{J}_3} = g_{J_1 J_2}^{J_3} \hat{g}_{\hat{J}_1 \hat{J}_2}^{\hat{J}_3} h_{J_1 \hat{J}_1 J_2 \hat{J}_2}^{J_3 \hat{J}_3} \quad (5.11)$$

where a careful calculation gives

$$h_{J_1 \hat{J}_1 J_2 \hat{J}_2}^{J_3 \hat{J}_3} = (-1)^{p\hat{p}+2p\hat{J}_3+2\hat{p}J_3} \prod_k \prod_{r, \hat{r} \in X_k} (r\sqrt{h/\pi} + \hat{r}\sqrt{\pi/h})^{\epsilon_k} \quad (5.12)$$

where the X'_k s are the direct product of intervals $[a_k, b_k] \otimes [\hat{a}_k, \hat{b}_k]$ and given by

$$X_k = \begin{cases} [1, 2J_1] \otimes [1, 2\hat{J}_1] \\ [1, 2J_1 - p] \otimes [1, 2\hat{J}_1 - \hat{p}] \\ [1, 2J_2] \otimes [1, 2\hat{J}_2] \\ [1, 2J_2 - p] \otimes [1, 2\hat{J}_2 - \hat{p}] \\ [2, 2J_3 + 1] \otimes [2, 2\hat{J}_3 + 1] \\ [2, 2J_3 + p + 1] \otimes [2, 2\hat{J}_3 + \hat{p} + 1] \\ [1, p] \otimes [1, \hat{p}] \end{cases}, \quad \text{with} \quad \epsilon_k = \begin{cases} +1 \\ -1 \\ +1 \\ -1 \\ +1 \\ -1 \\ -1 \end{cases}$$

and

$$p = J_1 + J_2 - J_3, \hat{p} = \hat{J}_1 + \hat{J}_2 - \hat{J}_3. \quad (5.13)$$

We note that G is symmetric under the exchange of 1 and 2. This is in agreement with the consistency of the fusion rules and the braiding.

Using the properties of the Γ function, the factor h can be absorbed partially or completely. A particular form which exhibits the symmetry between $(J_1, \hat{J}_1)$, $(J_2, \hat{J}_2)$ and $(-J_3 - 1, -\hat{J}_3 - 1)$ is

$$g_{J_1 \hat{J}_1 J_2 \hat{J}_2}^{J_3 \hat{J}_3} = (i/2)^{p+\hat{p}} \frac{H_{pp}(\hat{J}_1, \hat{J}_1) H_{pp}(\hat{J}_2, \hat{J}_2) H_{pp}(-J_3 - 1, -\hat{J}_3 - 1)}{H_{pp}(p/2, \hat{p}/2)} \quad (5.14)$$

with

$$H_{pp}(\hat{J}, \hat{J}) = \frac{\sqrt{\prod_{r=1}^p F(2\hat{J} + 1 + (2J - r + 1)h/\pi) \prod_{r=1}^{\hat{p}} F(2J + 1 + (2\hat{J} - \hat{r} + 1)\pi/h)}}{\prod_{r=1}^p \prod_{r=1}^{\hat{p}} [(2J - r + 1)\sqrt{h/\pi} + (2\hat{J} - \hat{r} + 1)\sqrt{\pi/h}]} \quad (5.15)$$

and $p = 1 + J_1 + J_2 + (-J_3 - 1)$ is symmetric.

The $H_{pp}(\hat{J}, \hat{J})$ function can be rewritten in a more useful form by absorbing all the denominator factors

$$H_{pp}(\hat{J}, \hat{J}) = e^{i\pi p\hat{p}/2} \prod_{\ell, \hat{\ell} \in \{\gamma\}} \left\{ F[R_\ell + (R_{\hat{\ell}} + (1 - \epsilon_{\hat{\ell}})/2)\pi/h](h/\pi)^{R_\ell - 2J + p - 1} \right\}^{\epsilon_{\hat{\ell}}/2} \\ \left\{ F[R_{\hat{\ell}} + (\hat{R}_\ell + (1 - \epsilon_\ell)/2)\pi/h](h/\pi)^{R_{\hat{\ell}} - 2\hat{J} + \hat{p} - 1} \right\}^{\epsilon_\ell/2} \quad (5.16)$$

where γ is an arbitrary path in the plane $(R, \hat{R})$ going from $(2J - p + 1, 2\hat{J} - \hat{p} + 1)$ to $(2J + 1, 2\hat{J} + 1)$ and

$\hat{\ell}$ labels the elementary path $(R_\ell, R_{\hat{\ell}}) \rightarrow (R_\ell, R_{\hat{\ell}} + \epsilon_{\hat{\ell}})$

ℓ labels the elementary path $(R_\ell, R_{\hat{\ell}}) \rightarrow (R_\ell + \epsilon_\ell, R_{\hat{\ell}})$

with $\epsilon_\ell, \epsilon_{\hat{\ell}} = 0, \pm 1$.

This allows to relate two G by a product of elementary shifts which have a simple form. From

$$\frac{H_{pp}(\hat{J} + 1/2, \hat{J})}{H_{pp}(\hat{J}, \hat{J})} = (h/\pi)^{-\hat{p}/2} \sqrt{\frac{F(2\hat{J} + 1 + (2J + 1)h/\pi)}{F(2\hat{J} - \hat{p} + 1 + (2J - p + 1)h/\pi)}} \quad (5.17)$$

$$\frac{H_{p+1\hat{p}}(\hat{J}, \hat{J})}{H_{pp}(\hat{J}, \hat{J})} = \sqrt{F(2\hat{J} - \hat{p} + 1 + (2J - p)h/\pi)} \quad (5.18)$$

$$\frac{H_{p+1\hat{p}}(p+1/2, \hat{p})}{H_{p\hat{p}}(J, \hat{J})} = (h/\pi)^{-\hat{p}/2} \sqrt{F(\hat{p}+1+(p+1)h/\pi)} \quad (5.19)$$

and related formula for $J \rightarrow J - 1/2$, $p \rightarrow \hat{p}$ we get

$$\frac{g_{J_1+1/2, \hat{J}_1, J_2-1/2, \hat{J}_2}^{J_3 \hat{J}_3}}{g_{J_1 \hat{J}_1 J_2 \hat{J}_2}^{J_3 \hat{J}_3}} = \sqrt{\frac{F(2\hat{J}_1+1+(2J_1+1)h/\pi)F(2\hat{J}_2-\hat{p}+1+(2J_2-p)h/\pi)}{F(2\hat{J}_2+1+2J_2h/\pi)F(2\hat{J}_1-\hat{p}+1+(2J_1-p+1)h/\pi)}}, \quad (5.20)$$

$$\frac{g_{J_1+1/2, \hat{J}_1, J_2+1/2, \hat{J}_2}^{J_3 \hat{J}_3}}{g_{J_1 \hat{J}_1 J_2 \hat{J}_2}^{J_3 \hat{J}_3}} = i/2(h\pi)^{-\hat{p}/2} \sqrt{F(2\hat{J}_1+1+(2J_1+1)h/\pi)} \times \sqrt{\frac{F(2\hat{J}_2+1+(2J_2+1)h/\pi)F(-2\hat{J}_3-\hat{p}-1-(2J_3+p+1)h/\pi)}{F(\hat{p}+1+(p+1)h/\pi)}}. \quad (5.21)$$

All other relations can be deduced by inversion of these relations or by the symmetry between J_i, h and $\hat{J}_i, \hat{h}$ or the symmetry between $J_1, J_2, -J_3 - 1$ (resp. $\hat{J}_i$).

6 CONCLUSION

As mentioned in the introduction, the earlier works only dealt with leading-order fusions, explicitly. In this case, one only retains the spin $J_1 + J_2$ in fusing operators with spins J_1 and J_2 . We began the present study, when we realized that, beyond this approximation, the OPA is not associative when it is considered too naively, that is at the level of primaries. This difficulty appears at two levels. First, in the product of three operators one sees that the dominant behaviour depends upon the way the three world-sheet points approach each other. Second, it is possible to define a limit such that this naive analysis gives associativity conditions over matrix-products of F and B matrices, but these relations are not satisfied. This fact is particularly clear for ξ fields. For them, OPE's at the level of primaries correspond to making q-tensor product of the representations, and this is notoriously non-associative. As a matter of fact, the 6-j symbols precisely encode this non-associativity (see the defining relation Eq.3.13), as it is well known¹⁸. Specifically, the naive analysis leads to Eq.3.12 without the fusing matrix F , or to Eq.3.13 without the 6-j symbol, which are clearly wrong relations. The general conclusion of the present work is that this difficulty is solved when the primaries are included using the compact formulae of the MS scheme, where all coefficients of the OPE are expressed by the V matrix element in Eqs.2.52, or 3.15. Then the coupling constant g_{JK}^L may be computed from the MS polynomial equations, starting from the particular case where one of the operators has $J = 1/2$.

¹⁸this is also true for ordinary groups.

As was pointed out several times[19, 20, 27], the $U_q(sl(2))$ quantum-group structure remarkably comes out of the holomorphic OPA of Liouville theory. It is interesting to think about a sort of reciprocal. Given the quantum group quantities – C.G. coefficients, 6-j symbols, universal R matrix –, the tensor product of representations defines a “multiplication” of representations which is not associative: this corresponds to the naive analysis mentioned above. Then, one may consider introducing additional quantum numbers such that the “product” becomes associative. A solution for this is given by the indices $\{\nu\}$ that characterize the descendants, and then the “product” becomes the OPE of holomorphic fields we have displayed. Indeed, the only way to have Eq.3.13 satisfied is that F be equal to a 6-j symbol. This forces the existence of the V ’s and fixes their OPE. Then some of the J ’s of the 6-j symbols have a natural interpretation as zero-modes which are shifted by the V fields as shown on Eq.A.12. It is likely that this is the unique possibility, although a proof of this fact would be beyond the scope of this article. In any case, our present study indicates that the relationship between $U_q(sl(2))$ and the holomorphic OPA of 2D gravity goes even more deeply than previously thought.

One may foresee future developments of the present work in several directions. The most interesting one at this point concerns the strong-coupling regime $1 < C < 25$, with complex h , where the present method is the only available so far. In ref.[20] a unitary-truncation theorem was derived for $C = 7, 13$, and 19 , by only considering leading-order OPE’s where it is sufficient to deal with primaries. This discussion may now be completed using the result of the present paper. Moreover, the solutions of the MS relations just displayed should allow us to check that the associated non-critical string theories are consistent as such, namely, that they satisfy duality relations between crossed channels similar to the ones of the Veneziano model. Another point is that, now that the full bootstrap is at hand, one may re-consider, with a much greater insight, the extension to negative J , handled in ref.[20] by means of a symmetry principle between spins J and $-J - 1$. Dealing with negative spins is unavoidable since they are the holomorphic components of exponentials of Liouville field with positive weights. These must be introduced for proper dressing by 2D gravity[27].

At a more ambitious level, understanding the connection between quantum group and 2D gravity more deeply is certainly a step toward unravelling the non-commutative geometry of the latter theory.

Acknowledgements

At an early stage of this work, one of us (J.-L.G.) was visiting the Mathematical Science Research Institute in Berkeley California. He is grateful for the generous financial support, the very stimulating surroundings and the warm hospitality from which he benefitted while participating in the program “Strings in Mathematics”.

This work was supported in part by the European Twinning Program, contrat #540022.

A APPENDIX A

In this appendix, some basic concepts of the present approach are recalled for completeness. The notations are identical to the ones used in the articles [19, 28, 20, 27]

to which we refer for details.

We recall some basic properties of the primary fields that came out as holomorphic components of exponentials of the Liouville field. Since one deals with functions of a single variable $\sigma - i\tau$ (see the beginning of part 2), one may work at $\tau = 0$ without loss of generality that is on the unit circle $u = e^{i\sigma}$. These holomorphic components may thus be regarded as functions of σ . The starting point is that, for trivial Verma modules, there exist two equivalent free fields:

$$\phi_j(\sigma) = q_0^{(j)} + p_0^{(j)}\sigma + i \sum_{n \neq 0} e^{-in\sigma} p_n^{(j)}/n, \quad j = 1, 2, \quad (\text{A.1})$$

such that (primes denote derivatives)

$$[\phi_1'(\sigma_1), \phi_1'(\sigma_2)] = [\phi_2'(\sigma_1), \phi_2'(\sigma_2)] = 2\pi i \delta'(\sigma_1 - \sigma_2), \quad p_0^{(1)} = -p_0^{(2)}, \quad (\text{A.2})$$

$$N^{(1)}(\phi_1')^2 + \phi_1''/\sqrt{\gamma} = N^{(2)}(\phi_2')^2 + \phi_2''/\sqrt{\gamma}. \quad (\text{A.3})$$

$N^{(1)}$ (resp. $N^{(2)}$) denote normal orderings with respect to the modes of ϕ_1 (resp. of ϕ_2). Eq.A.3 defines the stress-energy tensor and the coupling constant γ of the quantum theory. The former generates a representation of the Virasoro algebra with central charge $C = 3 + 1/\gamma$. At an intuitive level, the correspondence between ϕ_1 and ϕ_2 may be understood from the fact that the Verma modules, which they generate, coincide since the highest weights only depend upon $(p_0^{(1)})^2 = (p_0^{(2)})^2$. The chiral family is built up[9, 22, 23, 19] from the following operators

$$V_j = d_j N^{(j)}(e^{\sqrt{h/2\pi} \phi_j}), \quad \hat{V}_j = \hat{d}_j N^{(j)}(e^{\sqrt{\hat{h}/2\pi} \phi_j}), \quad j = 1, 2, \quad (\text{A.4})$$

$$h = \frac{\pi}{12} (C - 13 - \sqrt{(C - 25)(C - 1)}), \quad \hat{h} = \frac{\pi}{12} (C - 13 + \sqrt{(C - 25)(C - 1)}). \quad (\text{A.5})$$

In Eq.A.4, the zero modes are ordered, as is standard in string vertices, so that the operators are primary. Explicitly one has

$$N^{(j)}(e^{\sqrt{h/2\pi} \phi_j}) \equiv e^{\sqrt{h/2\pi} q_0^{(j)}} e^{\sqrt{h/2\pi} p_0^{(j)}} e^{-i\sigma h/4\pi} \times \\ \exp \left(\sqrt{h/2\pi} i \sum_{n < 0} e^{-in\sigma} p_n^{(j)}/n \right) \exp \left(\sqrt{h/2\pi} i \sum_{n > 0} e^{-in\sigma} p_n^{(j)}/n \right) \quad (\text{A.6})$$

with similar definitions for the hatted fields. The relation between h or $\hat{h}$ and C which is equivalent to

$$C = 1 + 6\left(\frac{h}{\pi} + \frac{\pi}{h} + 2\right) = 1 + 6\left(\frac{\hat{h}}{\pi} + \frac{\pi}{\hat{h}} + 2\right), \quad \text{with} \quad h\hat{h} = \pi^2, \quad (\text{A.7})$$

is such that V_j and $\hat{V}_j$ are solutions of the equations[21, 9]

$$-\frac{d^2 V_j(\sigma)}{d\sigma^2} + \left(\frac{h}{\pi}\right) \left(\sum_{n < 0} L_n e^{-in\sigma} + \frac{L_0}{2} + \left(\frac{h}{16\pi} - \frac{C-1}{24}\right) \right) V_j(\sigma) \\ + \left(\frac{\hat{h}}{\pi}\right) V_j(\sigma) \left(\sum_{n > 0} L_n e^{-in\sigma} + \frac{L_0}{2} \right) = 0 \quad (\text{A.8})$$

$$\begin{aligned}
& -\frac{d^2 \hat{V}_j(\sigma)}{d\sigma^2} + \left(\frac{\hat{h}}{\pi}\right) \left(\sum_{n<0} L_n e^{-in\sigma} + \frac{L_0}{2} + \left(\frac{h}{16\pi} - \frac{C-1}{24}\right)\right) \hat{V}_j(\sigma) \\
& + \left(\frac{\hat{h}}{\pi}\right) \hat{V}_j(\sigma) \left(\sum_{n>0} L_n e^{-in\sigma} + \frac{L_0}{2}\right) = 0
\end{aligned} \tag{A.9}$$

These are operator Schrödinger equations equivalent to the decoupling of Virasoro null-vectors[9, 22, 23]. Since there are two possible quantum modifications h and $\hat{h}$, there are four solutions. By operator product V_j , $j = 1, 2$, and $\hat{V}_j$, $j = 1, 2$, generate two infinite families of chiral fields $V_m^{(J)}$, $-J \leq m \leq J$, and $\hat{V}_m^{(\hat{J})}$, $-\hat{J} \leq \hat{m} \leq \hat{J}$; with $V_{-1/2}^{(1/2)} = V_1$, $V_{1/2}^{(1/2)} = V_2$, and $\hat{V}_{-1/2}^{(1/2)} = \hat{V}_1$, $\hat{V}_{1/2}^{(1/2)} = \hat{V}_2$. The fields $V_m^{(J)}$, $\hat{V}_m^{(\hat{J})}$, are of the type $(1, 2J+1)$ and $(2\hat{J}+1, 1)$, respectively, in the BPZ classification. For the zero-modes, it is simpler[19] to define the rescaled variables

$$\varpi = ip_0^{(1)} \sqrt{\frac{2\pi}{h}}; \quad \widehat{\varpi} = ip_0^{(1)} \sqrt{\frac{2\pi}{\hat{h}}}; \quad \widehat{\varpi} = \varpi \frac{h}{\pi}; \quad \varpi = \widehat{\varpi} \frac{\hat{h}}{\pi}. \tag{A.10}$$

The Hilbert space in which the operators ψ and $\hat{\psi}$ live, is a direct sum[19, 28, 20] of Verma modules $\mathcal{H}(\varpi)$. They are eigenstates of the quasi momentum ϖ , and satisfy $L_n|\varpi\rangle = 0$, $n > 0$; $(L_0 - \Delta(\varpi))|\varpi\rangle = 0$. The corresponding highest weights $\Delta(\varpi)$ may be rewritten as

$$\Delta(\varpi) \equiv \frac{1}{8\gamma} + \frac{(p_0^{(1)})^2}{2} = \frac{h}{4\pi} \left(1 + \frac{\pi}{h}\right)^2 - \frac{h}{4\pi} \varpi^2. \tag{A.11}$$

The commutation relations Eq.A.2 are to be supplemented by the zero-mode ones:

$$[q_0^{(1)}, p_0^{(1)}] = [q_0^{(2)}, p_0^{(2)}] = i.$$

It thus follows (see in particular Eq.A.6), that the fields V and $\hat{V}$ shift the quasi momentum $p_0^{(1)} = -p_0^{(2)}$ by a fixed amount. For an arbitrary c-number function f , one has

$$V_m^{(J)} f(\varpi) = f(\varpi + 2m) V_m^{(J)}, \quad \hat{V}_m^{(\hat{J})} f(\varpi) = f(\varpi + 2\hat{m}\pi/h) \hat{V}_m^{(\hat{J})}. \tag{A.12}$$

The fields V and $\hat{V}$ together with their products may be naturally restricted to discrete values of ϖ . They thus live in Hilbert spaces of the form¹⁹

$$\mathcal{H} \equiv \bigoplus_{n, \hat{n}=-\infty}^{+\infty} \mathcal{H}(\varpi_0 + n + \hat{n}\pi/h). \tag{A.13}$$

where $\mathcal{H}(\varpi_0 + n + \hat{n}\pi/h)$ are Verma modules. ϖ_0 is a constant. The $sl(2, C)$ -invariant vacuum corresponds to $\varpi_0 = 1 + \pi/h$, [19]. With this choice, we use the notation $\mathcal{H}_{J\hat{J}}$ instead of $\mathcal{H}(\varpi_0 + n + \hat{n}\pi/h)$. For h real, Eq.A.13 shows that the eigenvalues of ϖ may be most naturally assumed to be real. Thus $(p_0^{(1)})^\dagger = p_0^{(2)}$. As discussed several times[19], the natural hermiticity relation is $(\phi^{(1)}(\sigma))^\dagger = \phi^{(2)}(\sigma)$.

¹⁹ in the present paper, we only deal with V operators in detail, so that $\hat{n}$ is kept equal to zero

It follows from Eq.A.3 that this is consistent with the usual hermiticity relation $L_n^\dagger = L_{-n}$. As shown in [19], one has

$$(\psi_m^{(J)})^\dagger = \psi_{-m}^{(J)} \quad (\text{A.14})$$

The most general $(2\hat{J} + 1, 2J + 1)$ field $V_{m\hat{m}}^{(J\hat{J})} \sim V_m^{(J)} \hat{V}_{\hat{m}}^{(\hat{J})}$ has weight

$$\Delta_{J\hat{J}} = \frac{C-1}{24} - \frac{1}{24} \left((J + \hat{J} + 1)\sqrt{C-1} - (J - \hat{J})\sqrt{C-25} \right)^2 \quad (\text{A.15})$$

in agreement with Kac's formula. It is sometimes convenient to rewrite the operators $V_{M\hat{M}}^{(J\hat{J})}$ as $V^{\mu,\nu;\hat{\mu},\hat{\nu}}$ where

$$2J = \mu + \nu, \quad 2m = \nu - \mu, \quad 2\hat{J} = \hat{\mu} + \hat{\nu}, \quad 2\hat{m} = \hat{\nu} - \hat{\mu}. \quad (\text{A.16})$$

$V^{\mu,\nu;\hat{\mu},\hat{\nu}}(\sigma)$ is an operator whose normalization does not depend upon its indices: its only non-vanishing matrix element is

$$\langle \varpi | V^{\mu,\nu;\hat{\mu},\hat{\nu}}(\sigma) | \varpi - \mu + \nu + (-\hat{\mu} + \hat{\nu})\pi/h \rangle = e^{i\sigma(\varpi^2 - (\varpi - \mu + \nu + (-\hat{\mu} + \hat{\nu})\pi/h)^2)h/4\pi}. \quad (\text{A.17})$$

In Appendix E of ref.[19] fields $\psi_{m\hat{m}}^{(J\hat{J})}$ were defined by the relation

$$\psi_{m\hat{m}}^{(J\hat{J})}(\sigma) \equiv C^{\mu,\nu;\hat{\mu},\hat{\nu}} D^{\mu,\nu;\hat{\mu},\hat{\nu}}(\varpi) V^{\mu,\nu;\hat{\mu},\hat{\nu}}(\sigma), \quad (\text{A.18})$$

In the present article we use a condensed notation, by letting

$$E_{m\hat{m}}^{(J\hat{J})}(\varpi) \equiv C^{\mu,\nu;\hat{\mu},\hat{\nu}} D^{\mu,\nu;\hat{\mu},\hat{\nu}}(\varpi) \quad (\text{A.19})$$

where Eq.A.16 is to be used. According to Eqs.A.16 and E.3 of [19], $C^{\mu,\nu;\hat{\mu},\hat{\nu}}$ is given by

$$C^{\mu,\nu;\hat{\mu},\hat{\nu}} = C^{\mu,\nu} \hat{C}^{\hat{\mu},\hat{\nu}} \frac{\prod_{r=1}^{\mu+\nu} \prod_{\hat{r}=1}^{\hat{\mu}+\hat{\nu}} (r\sqrt{h/\pi} + \hat{r}\sqrt{\pi/h})}{\left[\prod_{s=1}^{\mu} \prod_{\hat{s}=1}^{\hat{\mu}} (s\sqrt{h/\pi} + \hat{s}\sqrt{\pi/h}) \prod_{t=1}^{\nu} \prod_{\hat{t}=1}^{\hat{\nu}} (t\sqrt{h/\pi} + \hat{t}\sqrt{\pi/h}) \right]}, \quad (\text{A.20})$$

$$C^{\mu,\nu} = \prod_{r=1}^{\mu+\nu} \Gamma(1 + rh/\pi) / \left[\prod_{s=1}^{\mu} \Gamma(1 + sh/\pi) \prod_{t=1}^{\nu} \Gamma(1 + th/\pi) \right], \quad (\text{A.21})$$

with similar formulae for $\hat{C}^{\hat{\mu},\hat{\nu}}$. The ϖ dependent part $D^{\mu,\nu;\hat{\mu},\hat{\nu}}(\varpi)$ is given²⁰ by Eqs.A.26a,b and E.4 of [19]:

$$D^{\mu,\nu;\hat{\mu},\hat{\nu}}(\varpi) = D^{\mu,\nu}(\varpi) \hat{D}^{\hat{\mu},\hat{\nu}}(\varpi) M^{\mu,\nu;\hat{\mu},\hat{\nu}}(\varpi) \quad (\text{A.22})$$

$$D^{\mu,\nu}(\varpi) = \prod_{t=1}^{\mu-\nu} \frac{\sqrt{\Gamma(-(\varpi - t + 1)h/\pi)}}{\sqrt{\Gamma((\varpi - t)h/\pi)}} \prod_{r=1}^{\mu} \Gamma((\varpi - r)h/\pi) \prod_{s=1}^{\nu} \Gamma((-\varpi - s)h/\pi), \quad (\text{A.23})$$

²⁰ up to a few misprints

if $\mu > \nu$, and

$$D^{\mu,\nu}(\varpi) = \prod_{t=1}^{\nu-\mu} \frac{\sqrt{\Gamma((\varpi+t-1)h/\pi)}}{\sqrt{\Gamma((-\varpi-t)h/\pi)}} \prod_{r=1}^{\mu} \Gamma((\varpi-r)h/\pi) \prod_{s=1}^{\nu} \Gamma((-\varpi-s)h/\pi), \quad (\text{A.24})$$

if $\nu > \mu$. Moreover, if for instance, $\mu > \nu$, $\hat{\mu} > \hat{\nu}$,

$$M^{\mu,\nu;\hat{\mu},\hat{\nu}}(\varpi) = \frac{\prod_{t=1}^{\mu-\nu} \prod_{\hat{t}=1}^{\hat{\mu}-\hat{\nu}} \sqrt{-(\varpi-t+1)\sqrt{h/\pi} + (\hat{t}-1)\sqrt{\pi/h}} \sqrt{(\varpi-t)\sqrt{h/\pi} - \hat{t}\sqrt{\pi/h}}}{\prod_{r=1}^{\mu} \prod_{\hat{r}=1}^{\hat{\mu}} ((\varpi-r)\sqrt{h/\pi} - \hat{r}\sqrt{\pi/h}) \prod_{s=1}^{\nu} \prod_{\hat{s}=1}^{\hat{\nu}} (-(\varpi+s)\sqrt{h/\pi} - \hat{s}\sqrt{\pi/h})}. \quad (\text{A.25})$$

Next, an other basis of operators, the ξ fiels were introduced in ref.[19] (see part 3 for the details). In [19, 20] their braiding algebra was completely determined. It was shown that it is given by the universal R-matrix of $U_q(sl(2))$ (see Eqs.3.21 - 3.23).

B APPENDIX B

In this section we comment about the z -dependence of the fusion equations derived in section 2, and connect it with the standard form where translation invariance is manifest. First consider Eq.2.15. It follows from Eq.2.59 hat the states $|\varpi_1 >$, and $< \varpi_4|$ are obtained from the limits

$$\lim_{z_1 \rightarrow 0} V_{-J_1}^{(J_1)}(z_1) |\varpi_0 > = |\varpi_1 >, \quad \lim_{z_4 \rightarrow \infty} (-z_4)^{2\Delta_4} < -\varpi_0 | V_{J_4+1,+1}^{(-J_4-1,-1)}(z_4) = < \varpi_4|. \quad (\text{B.1})$$

Thus we have

$$\begin{aligned} < \varpi_4 | V_{\pm 1/2}^{(1/2)}(z_3) V_{m_2}^{(J_2)}(z_2) | \varpi_1 > = \\ \lim_{z_4 \rightarrow \infty, z_1 \rightarrow 0} (-z_4)^{2\Delta_4} < -\varpi_0 | V_{J_4+1,1}^{(-J_4-1,-1)}(z_4) V_{\pm 1/2}^{(1/2)}(z_3) V_{m_2}^{(J_2)}(z_2) V_{-J_1}^{(J_1)}(z_1) | \varpi_0 > . \end{aligned} \quad (\text{B.2})$$

Similarly one sees that

$$\begin{aligned} < \varpi_1 | V_{m_4}^{(J_4)}(z'_4) V_{\pm 1/2}^{(1/2)}(z'_3) | \varpi_2 > = \\ \lim_{z'_1 \rightarrow \infty, z'_2 \rightarrow 0} (-z'_1)^{2\Delta_1} < -\varpi_0 | V_{J_1+1,1}^{(-J_1-1,-1)}(z'_4) V_{m_4}^{(J_4)}(z'_4) V_{\pm 1/2}^{(1/2)}(z'_3) V_{-J_2}^{(J_2)}(z'_2) | \varpi_0 > . \end{aligned} \quad (\text{B.3})$$

Since $|\varpi_0 >$ and $< -\varpi_0|$ are the right and left $sl(2, C)$ -invariant states, respectively, it follows that any matrix element of primary fields $< -\varpi_0 | A_1 A_2 A_3 A_4 | \varpi_0 >$, is of the form $f / \prod_{i,j} (z_i - z_j)^{\Delta_i + \Delta_j - \Delta}$, where f is invariant²¹ under projective (Möbius) transformations of the z 's, and $\Delta = \sum_j \Delta_j / 3$. It is thus clear that the above z and z' variables should be related by a projective transformation. This is verified as follows. Eq.2.15 is such that $z'_4 = z_2$. Thus the projective transformation is characterized by the condition $(\infty, z_2, 0) \rightarrow (z_2, 0, \infty)$, and one has

$$z'_k = z_2(1 - z_2/z_k). \quad (\text{B.4})$$

²¹it may be regarded as a function of the anharmonic (cross) ratio of the z 's

This immediately gives $z'_3 = z_2(1 - z_2/z_3)$ in agreement with Eq.2.15. Moreover there is the Jacobian factor $J = \prod_k (dz'_k/dz_k)^{\Delta_k}$ which arises due to the transformation law Eq.2.6 of the primary fields. Since $dz'/dz = (z_2/z)^2$, one has

$$J = z_2^{2(\Delta_1 + \Delta_3 + \Delta_4)} \prod_{l=1,3,4} z_l^{-2\Delta_l} \sim_{z_1 \rightarrow 0} (-z'_1)^{2\Delta_1} (z_4)^{-2\Delta_4} z_2^{2(\Delta_3 + \Delta_4 - \Delta_1)} z_3^{-2\Delta_3}$$

The first two terms on the right-hand side coincide with the factors in Eqs.B.2, and B.3 that respectively ensure the finiteness of the limits $z_4 \rightarrow \infty$, and $z'_1 \rightarrow \infty$. The other factors are equal to the first two terms of the right-hand side of Eq.2.15.

Finally, we show how invariance under z translations may be recovered. Assume, as usual that Eq.2.22 is true for arbitrary matrix element. Then we may write

$$\begin{aligned} V_{\pm 1/2}^{(1/2)}(z_3) V_{m_2}^{(J_2)}(z_2) &= \left(\frac{z_2}{z_3}\right)^{2\Delta_{1/2}} \sum_{\epsilon=\pm 1} f_{\pm 1, \epsilon}(J_2, m_2; \varpi) \times \\ &\sum_{\{\nu\}} V_{m_2 \pm 1/2}^{(J_\epsilon, \{\nu\})}(z_2) < \varpi_\epsilon, \{\nu\} | V_{-\epsilon/2}^{(1/2)} \left(z_2 \left(1 - \frac{z_2}{z_3}\right) \right) | \varpi_2 >, \end{aligned} \quad (\text{B.5})$$

where ϖ is the zero-mode operator. This formula is invariant under the transformation $z \rightarrow \lambda z$ which preserves the points 0 and ∞ associated with the matrix element we started from. Covariance under z translations ($z \rightarrow z + a$), is achieved by performing the Möbius transformation $z \rightarrow \zeta = -z_2 z_3 / z$ on the three operators $V_{\pm 1/2}^{(1/2)}(z_3)$, $V_{m_2}^{(J_2)}(z_2)$, $V_{m_2 \pm 1/2}^{(J_\epsilon, \{\nu\})}(z_2)$, and rescaling the argument of $V_{-\epsilon/2}^{(1/2)}$ in the matrix element so that $z_2(1 - z_2/z_3) \rightarrow z_3 - z_2$. Taking account of the transformation factor of Eq.2.6, one finds

$$\begin{aligned} V_{\pm 1/2}^{(1/2)}(-z_2) V_{m_2}^{(J_2)}(-z_3) &= \sum_{\epsilon=\pm 1} f_{\pm 1, \epsilon}(J_2, m_2; \varpi) \times \\ &\sum_{\{\nu\}} V_{m_2 \pm 1/2}^{(J_\epsilon, \{\nu\})}(-z_3) < \varpi_\epsilon, \{\nu\} | V_{-\epsilon/2}^{(1/2)}(z_3 - z_2) | \varpi_2 >. \end{aligned} \quad (\text{B.6})$$

Up to a change of sign in the definition of the z variable, this takes the usual form.

C APPENDIX C

This section is devoted to the explicit derivation of Eq. 3.8, that is

$$\begin{aligned} \sum_{\eta, \mu, (\eta + \mu = n)} |1/2, \varpi)_\alpha^\eta | J, \varpi + 2\eta)_M^\mu N_{\eta}^{1/2 J; J + \epsilon/2; \varpi} \Big|_\mu^n = \\ (1/2, \alpha; J, M | J + \epsilon/2, \varpi)_{M+\alpha}^n \end{aligned} \quad (\text{C.1})$$

The case $\epsilon = 1$ is already discussed in ref.[19] where the general fusion coefficients to leading order were determined. We shall, nevertheless rediscuss this case first for completeness, and since the case $\epsilon = -1$ is exactly similar. Letting, $n = m + 1/2$, one sees that we have to deal with the expression

$$|1/2, \varpi)_\alpha^{-1/2} | J, \varpi - 1)_M^m \frac{[\varpi + J + m]}{[\varpi]} + |1/2, \varpi)_\alpha^{1/2} | J, \varpi + 1)_M^{m-1} \frac{[\varpi - J + m - 1]}{[\varpi]}. \quad (\text{C.2})$$

The first thing is to get rid of the $[\varpi]$ denominators. This is done by using the following recurrence relation satisfied by the coefficients $|J, \varpi)_M^m$:

$$|J, \varpi + 1)_M^{m-1} = e^{-ih/2} \frac{[\varpi + J + m]}{[J - m + 1]} |J, \varpi - 1)_M^m - e^{ih(M-1/2)} \frac{[\varpi]}{[J - m + 1]} |J, \varpi)_M^m. \quad (\text{C.3})$$

This relation was established in ref.[19] in order to derive the braiding matrix of the ξ fields (Eqs.A.35-39) from the one of the ψ fields.

Next, we proceed by only considering Eq.C.2 for $\alpha = -1/2$, since this other value is treated in exactly the same way. Let us transform this relation by using the explicit expressions for $|1/2, \varpi)_\alpha^{\pm 1/2}$, that is,

$$|1/2, \varpi)_{\pm 1/2}^{\pm 1/2} = e^{ih(\varpi \pm 1/2)}, \quad |1/2, \varpi)_{\pm 1/2}^{\mp 1/2} = e^{-ih\varpi/2}. \quad (\text{C.4})$$

One gets

$$\frac{e^{-ih(\varpi+1)/2}}{[J - m + 1]} \left\{ e^{ih(J-m+1)} [\varpi + J + m] |J, \varpi - 1)_M^m - e^{ihM} [\varpi - J + m - 1] |J, \varpi)_M^m \right\}.$$

Eq.A.28 shows that the coefficients $|J, \varpi)_M^m$ are of the form $\sum_t e^{iht(\varpi+m)} A_M^t(J, m)$. Accordingly, the Fourier expansion of the last equation is

$$\frac{e^{-ih(\varpi+1)/2}}{[J - m + 1]} \sum_t e^{iht(\varpi+m)} e^{ih(M+J+1-t-m)/2} \left\{ \left[\frac{3 + 3J - M - t - m}{2} \right] A_M^{t-1}(J, m) + \left[\frac{1 + J + M + t + m}{2} \right] A_M^{t+1}(J, m) \right\}.$$

Using the explicit expression of $A_M^t(J, m)$, one sees that this last expression is equivalent to

$$e^{ih(-\varpi+M+J-m+1/2)/2} \sqrt{\frac{[J - M + 1]}{[2J + 1]}} \times \sum_t e^{iht(\varpi+m-1/2)} A_{M-1/2}^{t-1/2}(J + 1/2, m - 1/2).$$

Thus we finally get

$$\begin{aligned} |1/2, \varpi)_{-1/2}^{-1/2} |J, \varpi - 1)_M^m \frac{[\varpi + J + m]}{[\varpi]} + |1/2, \varpi)_{-1/2}^{1/2} |J, \varpi + 1)_M^{m-1} \frac{[\varpi - J + m - 1]}{[\varpi]} \\ = \sqrt{\frac{[J - M + 1]}{[2J + 1]}} e^{ih(J+M)/2} |J + 1/2, \varpi)_{M-1/2}^{m-1/2}. \end{aligned} \quad (\text{C.5})$$

This is the desired relation since (see, e.g. ref.[19])

$$(1/2, -1/2; J, M | J + 1/2) = e^{ih(J+M)/2} \sqrt{\frac{[J - M + 1]}{[2J + 1]}}.$$

Let us next turn to the case $\epsilon = -1$. Instead of Eq.C.2, one has to deal with

$$\frac{i}{2 \sin(h) \sqrt{[2J][2J + 1]}} \left\{ |1/2, \varpi)_\alpha^{-1/2} |J, \varpi - 1)_M^m \frac{[J + m]}{[\varpi]} \right.$$

$$+|1/2, \varpi)_\alpha^{1/2}|J, \varpi + 1)_M^{m-1} \frac{\lfloor -J + m - 1 \rfloor}{\lfloor \varpi \rfloor} \Big\}. \quad (\text{C.6})$$

Again, we consider the case $\alpha = -1/2$, and make use of Eq.C.3. The last expression becomes

$$\frac{-ie^{-ih(\varpi+1)/2}}{2 \sin(h) \sqrt{\lfloor 2J \rfloor \lfloor 2J + 1 \rfloor}} \left\{ e^{-ih(J+m)} |J, \varpi - 1)_M^m - e^{ihM} |J, \varpi)_M^m \right\}.$$

Its Fourier expansion is found to be

$$-\frac{e^{-ih(\varpi+1)/2}}{\sqrt{\lfloor 2J \rfloor \lfloor 2J + 1 \rfloor}} \sum_t e^{iht(\varpi+m)} e^{ih(M-J-t-m)/2} \left\lfloor \frac{J+M+t+m}{2} \right\rfloor A_M^t(J, m).$$

Using the explicit expression for $A_M^t(J, m)$, one verifies finally that this leads to

$$\begin{aligned} & \frac{i}{2 \sin(h) \sqrt{\lfloor 2J \rfloor \lfloor 2J + 1 \rfloor}} \left\{ |1/2, \varpi)_\alpha^{-1/2} |J, \varpi - 1)_M^m \frac{\lfloor J + m \rfloor}{\lfloor \varpi \rfloor} \right. \\ & \quad \left. + |1/2, \varpi)_\alpha^{1/2} |J, \varpi + 1)_M^{m-1} \frac{\lfloor -J + m - 1 \rfloor}{\lfloor \varpi \rfloor} \right\} \\ & = -e^{ih(-J+M-1/2)/2} \sqrt{\frac{\lfloor J + M \rfloor}{\lfloor 2J + 1 \rfloor}} |J - 1/2, \varpi)_{M+1/2}^{m-1/2}. \end{aligned} \quad (\text{C.7})$$

This is the desired result since

$$(1/2, -1/2; J, M | J - 1/2) = -e^{ih(-J+M-1)/2} \sqrt{\frac{\lfloor J + M \rfloor}{\lfloor 2J + 1 \rfloor}}$$

One may check the case $\alpha = 1/2$ in the same way, using the fact that

$$\begin{aligned} (1/2, 1/2; J, M | J + 1/2) &= e^{-ih(J-M)/2} \sqrt{\frac{\lfloor J + M + 1 \rfloor}{\lfloor 2J + 1 \rfloor}} \\ (1/2, 1/2; J, M | J - 1/2) &= e^{ih(J+M+1)/2} \sqrt{\frac{\lfloor J - M \rfloor}{\lfloor 2J + 1 \rfloor}} \end{aligned} \quad (\text{C.8})$$

References

- [1] E. Verlinde, *Nucl. Phys.* **B300** (1988) 360.
- [2] A. Tsuchia, Y. Kanie, *Advanced Study in Pure Mathematics*, **16**, (1987), 297.
- [3] T. Khono, *Ann. Inst. Fourier* (Grenoble) **37**, **N:4**, (1987), 139.
- [4] J. Fröhlich, 1987 Cargèse Lectures, in *Non-Perturbative Quantum Field Theory*, G. 't Hooft editor, Plenum Press (1988).
- [5] G. Felder, J. Fröhlich, J. Keller, *Comm. Math. Phys.* **124** (1989) 646.

- [6] J. Fröhlich, C. King, *Int. J. Mod. Phys.* **A4** (1989) 5321.
- [7] K.H. Rehren, *Comm. Math. Phys.* **116** (1988) 675.
- [8] K.H. Rehren, B. Schroer, *Nucl. Phys.* **B312** (1989) 715.
- [9] J.-L. Gervais, A. Neveu, *Nucl. Phys.* **B238** (1984) 125,
Nucl. Phys. **B238** (1984) 396.
- [10] G. Moore, N. Seiberg, *Comm. Math. Phys.* **123** (1989) 77.
- [11] G. Moore, N. Reshetikhin, *Nucl. Phys.* **B238** (1989) 557.
- [12] V. Pasquier, *Comm. Math. Phys.* **118** (1988) 355.
- [13] O. Babelon, *Phys. Lett.* **B215** (1988) 523.
- [14] L. Alvarez-Gaumé, C. Gomez, G. Sierra, *Nucl. Phys.* **B319** (1989. 155) *Phys. Lett.* **B220** (1989) 142, *Nucl. Phys.* **B[FS]** (1990) 347.
- [15] E. Witten, *Nucl. Phys.* **B330** (1990) 205.
- [16] I. Todorov, *Proceedings of the VIII Clausthal-Zellerfeld Workshop on Mathematical Physics* Springer Lecture Notes, Springer; L. Hadjivanov, R. Paunov, I. Todorov, *Nucl. Phys.* **B356** (1991) 387.
- [17] P. Furlan, A. Ganchev, V. Petkova, *Nucl. Phys.* **B343** (1990) 205.
- [18] K. Gawędzki, *Nucl. Phys. (Proc. Suppl.)* **18B** (1990) 78.
- [19] J.-L. Gervais, *Comm. Math. Phys.* **130** (1990) 257.
- [20] J.-L. Gervais, *Comm. Math. Phys.* **138** (1991) 301.
- [21] J.-L. Gervais, A. Neveu, *Nucl. Phys.* **B224** (1983) 329.
- [22] J.-L. Gervais, A. Neveu, *Nucl. Phys.* **B257[FS14]** (1985) 59.
- [23] J.-L. Gervais, A. Neveu, *Nucl. Phys.* **B264** (1986) 557.
- [24] C. Gomez, G. Sierra, *Nucl. Phys.* **B352** (1991) 791.
- [25] A. Kirilov, N. Reshetikhin, *Infinite Dimensional Lie Algebras and Groups, Advanced Study in Mathematical Physics* vol. **7**, Proceedings of the 1988 Marseille Conference, V. Kac editor, p. 285, World scientific.
- [26] E. Cremmer, J.-L. Gervais, *Comm. Math. Phys.* **144** (1992) 279.
- [27] J.-L. Gervais, “Quantum group derivation of 2D gravity-matter coupling” Invited talk at the Stony Brook meeting *String and Symmetry 1991*. “Gravity-Matter couplings from Liouville Theory”, preprint LPTENS 91-22, hep-th/9205034, *Nucl. Phys.* to be published.
- [28] J.-L. Gervais, *Phys. Lett.* **B243** (1990) 85.

- [29] G. Ponzano, T. Regge, in “Spectroscopic and group theoretical methods in Physics”, F. Bloch ed., North Holland, Amsterdam, 1968. B. Hasslacher, M. Perry, *Phys. Lett.* **103B** (1981) 21, S. M. Lewis, *Phys. Lett.* **122B** (1983) 265.
- [30] O. Babelon, *Comm. Math. Phys.* **139** (1991) 619.
- [31] J.-L. Gervais, B. Sakita, *Nucl. Phys.* **B34** (1971) 477.
- [32] Bo-yu Hou, Bo-yuan Hou, Zhong-qi Ma, *Comm. Theor. Phys.*, **13**, 181, (1990); *Comm. Theor. Phys.*, **13**, 1990, 341.